



Antibodies

11-308-C025

## Monoclonal Antibody to CD27 Purified Antibody (0.025 mg)

|                             |   |
|-----------------------------|---|
| <b>Clone:</b>               | LT27  |
| <b>Isotype:</b>             | Mouse IgG2a   |
| <b>Specificity:</b>         | The antibody LT27 reacts with CD27 (T14), a 50-55 kDa type I transmembrane glycoprotein (member of the TNF-receptor superfamily) expressed on medullary thymocytes, peripheral T lymphocytes, some B lymphocytes and NK cells.<br>HLDA V; WS Code T T-CD27.01   |
| <b>Regulatory Status:</b>   | RUO   |
| <b>Immunogen:</b>           | Human peripheral blood lymphocytes  |
| <b>Species Reactivity:</b>  | Human   |
| <b>Application:</b>         | Flow Cytometry<br>Recommended dilution: 2 µg/ml   |
| <b>Purity:</b>              | > 95% (by SDS-PAGE)   |
| <b>Purification:</b>        | Purified by precipitation and chromatography  |
| <b>Concentration:</b>       | 1 mg/ml   |
| <b>Storage Buffer:</b>      | Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.4   |
| <b>Storage / Stability:</b> | Store at 2-8°C. Do not freeze. Do not use after expiration date stamped on vial label.  |
| <b>Expiration:</b>          | See vial label  |
| <b>Lot Number:</b>          | See vial label  |
| <b>Background:</b>          | CD27 is a transmembrane 55 kDa protein of the nerve growth factor-receptor family, expressed as a disulfide-linked homodimer on mature thymocytes, peripheral blood T cells and a subpopulation of B cells. Activation of T cells via TCR-CD3 complex results in upregulation of CD27 expression on the plasma membrane as well as in the release of its soluble 28-32 kDa form, sCD27, detected in the plasma, urine or spinal fluid. This sCD27 is an important prognostic marker of acute and chronic B cell malignancies. RgpA, a cysteine proteinase, although activating T cells through the protease-activated receptors (PARs), degrades CD27 and counteracts T cell activation mediated by CD27 and its ligand CD70. |

**For laboratory research only, not for drug, diagnostic or other use.**



**Antibodies**

- References:**
- \*van Oers MH, Pals ST, Evers LM, van der Schoot CE, Koopman G, Bonfrer JM, Hintzen RQ, von dem Borne AE, van Lier RA: Expression and release of CD27 in human B-cell malignancies. *Blood*. 1993 Dec 1;82(11):3430-6.
  - \*Molica S, Vitelli G, Levato D, Crispino G, Dell'Olio M, Dattilo A, Matera R, Gandolfo GM, Musto P. CD27 in B-cell chronic lymphocytic leukemia. Cellular expression, serum release and correlation with other soluble molecules belonging to nerve growth factor receptors (NGFr) superfamily. *Haematologica*. 1998 May;83(5):398-402.
  - \*Kara IO, Sahin B, Gunesacar R: Expression of soluble CD27 and interleukins-8 and -10 in B-cell chronic lymphocytic leukemia: correlation with disease stage and prognosis. *Adv Ther*. 2007 Jan-Feb;24(1):29-40.
  - \*Yun LW, Decarlo AA, Hunter N: Blockade of protease-activated receptors on T cells correlates with altered proteolysis of CD27 by gingipains of *Porphyromonas gingivalis*. *Clin Exp Immunol*. 2007 Nov;150(2):217-29.
  - \*Leukocyte Typing V., Schlossman S. et al. (Eds.), Oxford University Press (1995).
  - \*Kanderova V, Kuzilkova D, Stuchly J, Vaskova M, Brdicka T, Fiser K, Hrusak O, Lund-Johansen F, Kalina T: High-resolution Antibody Array Analysis of Childhood Acute Leukemia Cells. *Mol Cell Proteomics*. 2016 Apr;15(4):1246-61.

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at [www.exbio.cz](http://www.exbio.cz).

**For laboratory research only, not for drug, diagnostic or other use.**

---

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic  
Tel: +420 261 090 666 | Fax: +420 261 090 660 | [orders@exbio.cz](mailto:orders@exbio.cz) | [www.exbio.cz](http://www.exbio.cz)