



11-403-C100

Monoclonal Antibody to CD261 / TRAIL-R1 Purified Antibody (0.1 mg)

Clone:	DR-4-02
Isotype:	Mouse IgG1
Specificity:	The antibody DR-4-02 recognizes TRAIL-R1 (DR4), a human death receptor 4 (468 amino acids) expressed in most human tissues (spleen, peripheral blood leucocytes, thymus) and in a variety of tumour-derived cell lines.
Regulatory Status:	RUO
Immunogen:	Fusion protein containing the extracellular part of TRAIL-R1 and the constant part of the heavy chain of the human IgG1.
Species Reactivity:	Human
Application:	Flow Cytometry Recommended dilution: 3-5 µg/ml Immunoprecipitation Immunocytochemistry Functional Application Soluble antibody DR-4-02 blocks apoptosis triggered by a ligand (TRAIL). Plastic-immobilized (cross-linked) DR-4-02 antibody induces apoptosis in sensitive cells. Recommended dilution of antibody: 2-3 µg/ml in cultivation medium Final concentration of TRAIL: 20-200 ng/ml Application note: It is recommended to add the antibody 15 min before addition of TRAIL.
Purity:	> 95% (by SDS-PAGE)
Purification:	Purified by protein-A affinity chromatography
Concentration:	1 mg/ml
Storage Buffer:	Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.4
Storage / Stability:	Store at 2-8°C. Do not freeze. Do not use after expiration date stamped on vial label.
Expiration:	See vial label
Lot Number:	See vial label
Background:	TRAIL-R1 (CD261, DR4) is a type I transmembrane protein, also called TRAIL receptor 1. The ligand for this DR4 death receptor has been identified and termed TRAIL, which is a member of the TNF family. DR4, as many other receptors (Fas, TNFR1, etc.), mediates apoptosis and NF kappaB activation in many cells and tissues. Apoptosis, a programmed cell death, is a operating process in normal cellular differentiation and development of multicellular organisms. Apoptosis is induced by coupled of certain cytokines (TNF family - TNF, Fas ligand) and their death domain containing receptors (TNFR1, Fas receptor).

For laboratory research only, not for drug, diagnostic or other use.



Antibodies

- References:**
- *Corallini F, Milani D, Nicolini V, Secchiero P: TRAIL, caspases and maturation of normal and leukemic myeloid precursors. *Leuk Lymphoma*. 2006 Aug;47(8):1459-68.
 - *Simova S, Klima M, Cermak L, Sourkova V, Andera L: Arf and Rho GAP adapter protein ARAP1 participates in the mobilization of TRAIL-R1/DR4 to the plasma membrane. *Apoptosis*. 2008 Mar;13(3):423-36.

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at www.exbio.cz.

For laboratory research only, not for drug, diagnostic or other use.

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic
Tel: +420 261 090 666 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz