



11-449-C100

Monoclonal Antibody to CD62L Purified Antibody (0.1 mg)

Clone:	LT-TD180
Isotype:	Mouse IgG1
Specificity:	The antibody LT-TD180 reacts with CD62L (L-selectin), a 74-95 kDa single chain type I glycoprotein expressed on most peripheral blood B lymphocytes, T lymphocytes, monocytes and granulocytes; it is also present on a subset of NK cells and certain hematopoietic malignant cells.
Regulatory Status:	RUO
Immunogen:	Peripheral blood leukocytes
Species Reactivity:	Human
Application:	Flow Cytometry
Purity:	> 95% (by SDS-PAGE)
Purification:	Purified by precipitation and chromatography
Concentration:	1 mg/ml
Storage Buffer:	Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.4
Storage / Stability:	Store at 2-8°C. Do not freeze. Do not use after expiration date stamped on vial label.
Expiration:	See vial label
Lot Number:	See vial label
Background:	CD62L (L-selectin) is an adhesion glycoprotein that is constitutively expressed on the cell surface of leukocytes and mediates their homing to inflammatory sites and peripheral lymph nodes by enabling rolling along the venular wall. CD62L is also involved in activation-induced neutrophil aggregation. Activation-dependent CD62L shedding, however, counteracts neutrophil rolling. CD62L has also signaling roles including enhance of chemokine receptor expression. Similarly to CD62P, the major ligand of CD62L is PSGL-1 (P-selectin glycoprotein ligand-1).

For laboratory research only, not for drug, diagnostic or other use.

**Antibodies****References:**

- *Von Andrian UH, Hansell P, Chambers JD, Berger EM, Torres Filho I, Butcher EC, Arfors KE: L-selectin function is required for beta 2-integrin-mediated neutrophil adhesion at physiological shear rates in vivo. *Am J Physiol.* 1992 Oct;263(4 Pt 2):H1034-44.
- *von Andrian UH, Chambers JD, Berg EL, Michie SA, Brown DA, Karolak D, Ramezani L, Berger EM, Arfors KE, Butcher EC. L-selectin mediates neutrophil rolling in inflamed venules through sialyl LewisX-dependent and -independent recognition pathways. *Blood.* 1993 Jul 1;82(1):182-91.
- *Simon SI, Burns AR, Taylor AD, Gopalan PK, Lynam EB, Sklar LA, Smith CW: L-selectin (CD62L) cross-linking signals neutrophil adhesive functions via the Mac-1 (CD11b/CD18) beta 2-integrin. *J Immunol.* 1995 Aug 1;155(3):1502-14.
- *Ding Z, Issekutz TB, Downey GP, Waddell TK: L-selectin stimulation enhances functional expression of surface CXCR4 in lymphocytes: implications for cellular activation during adhesion and migration. *Blood.* 2003 Jun 1;101(11):4245-52.
- *Ramachandran V, Williams M, Yago T, Schmidtke DW, McEver RP: Dynamic alterations of membrane tethers stabilize leukocyte rolling on P-selectin. *Proc Natl Acad Sci U S A.* 2004 Sep 14;101(37):13519-24.
- *Kuttruff S, Koch S, Kelp A, Pawelec G, Rammensee HG, Steinle A: NKp80 defines and stimulates a reactive subset of CD8 T cells. *Blood.* 2009 Jan 8;113(2):358-69.
- *Linnebacher M, Wienck A, Boeck I, Klar E: Identification of an MSI-H tumor-specific cytotoxic T cell epitope generated by the (-1) frame of U79260(FTO). *J Biomed Biotechnol.* 2010;2010:841451.
- *Simone R, Zicca A, Saverino D: The frequency of regulatory CD3+CD8+CD28-CD25+ T lymphocytes in human peripheral blood increases with age. *J Leukoc Biol.* 2008 Dec;84(6):1454-61.

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at www.exbio.cz.

For laboratory research only, not for drug, diagnostic or other use.

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic
Tel: +420 261 090 666 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz