



11-461-C025

Monoclonal Antibody to CD262 / TRAIL-R2 Purified Antibody (0.025 mg)

| | |
|-----------------------------|--|
| Clone: | DR5-01-1 |
| Isotype: | Mouse IgG1 |
| Specificity: | The antibody DR5-01-1 recognizes an extracellular domain of TRAIL-R2 (DR5). TRAIL-R2 is one of two TNF superfamily member intracellular death domain containing receptors for TRAIL (APO2L). |
| Regulatory Status: | RUO |
| Immunogen: | Recombinant fusion protein of human IgG heavy chain and extracellular domain of DR5. |
| Species Reactivity: | Human |
| Application: | Flow Cytometry Recommended dilution: 1-10 µg/ml Positive control: JURKAT human peripheral blood leukemia T cell line |
| Purity: | > 95% (by SDS-PAGE) |
| Purification: | Purified by protein-A affinity chromatography |
| Concentration: | 1 mg/ml |
| Storage Buffer: | Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.4 |
| Storage / Stability: | Store at 2-8°C. Do not freeze. Do not use after expiration date stamped on vial label. |
| Expiration: | See vial label |
| Lot Number: | See vial label |
| Background: | TRAIL-R2 (CD262, DR5) is one of two TNF superfamily member intracellular death domain containing receptors for TRAIL (APO2L). Apoptosis, or programmed cell death, occurs during normal cellular differentiation and development of multicellular organisms. Apoptosis is induced by certain cytokines including tumor necrosis factor (TNF) and Fas ligand in the TNF family through their death domain containing receptors, TNF receptor 1 (TNFR1) and Fas, respectively. Another member in the TNF family has been identified and designated TRAIL (for TNF related apoptosis inducing ligand) and Apo2L (for Apo2 ligand). Receptors for TRAIL include two death domain containing receptors, DR4 and DR5, as well as two decoy receptors, DcR1 and DcR2, lacking the intracellular signaling death domain. DcR1 (also called TRID), like the related death receptors DR4 and DR5, contains two extracellular cysteine rich domains. However, DcR1 contains no intracellular death domain and is thus incapable of signaling apoptosis. It has been suggested DcR1 is responsible for TRAIL resistance in normal human tissues including heart, placenta, lung, liver, kidney, spleen, and bone marrow. DR5 is a member of the TNF receptor superfamily, and contains an intracellular death domain. This receptor can be activated by tumor necrosis factor related apoptosis inducing ligand (TNFSF10/TRAIL/APO2L), and transduces apoptosis signal. Studies with FADD deficient mice suggested that FADD, a death domain containing adaptor protein, is required for the apoptosis mediated by this protein. |

For laboratory research only, not for drug, diagnostic or other use.



Antibodies

References: *Corallini F, Milani D, Nicolini V, Secchiero P: TRAIL, caspases and maturation of normal and leukemic myeloid precursors. *Leuk Lymphoma*. 2006 Aug;47(8):1459-68.

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's terms and conditions which are available at www.exbio.cz.

For laboratory research only, not for drug, diagnostic or other use.

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic
Tel: +420 261 090 666 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz