



11-640-C100

Monoclonal Antibody to CD209 Purified Antibody (0.1 mg)

Clone:	UW60.1
Isotype:	Mouse IgG1
Specificity:	The mouse monoclonal antibody UW60.1 recognizes human CD209 (DC-SIGN), a 44 kDa transmembrane receptor expressed on the surface of dendritic cells and macrophages.
Regulatory Status:	RUO
Immunogen:	CD209-His-tagged fusion protein
Species Reactivity:	Human
Application:	Flow Cytometry Immunohistochemistry Immunocytochemistry
Purity:	> 95% (by SDS-PAGE)
Purification:	Purified by protein-A affinity chromatography
Concentration:	1 mg/ml
Storage Buffer:	Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.4
Storage / Stability:	Store at 2-8°C. Do not freeze. Do not use after expiration date stamped on vial label.
Expiration:	See vial label
Lot Number:	See vial label
Background:	CD209, also known as DC-SIGN (dendritic cell-specific ICAM-3-grabbing nonintegrin) is a transmembrane receptor expressed on the surface of dendritic cells and macrophages, which recognizes numerous pathogens ranging from parasites to viruses. Its N-terminal domain is transmembrane, whereas a tandem-repeat neck domain and the C terminal C-type lectin carbohydrate recognition domain have dual function as a pathogen recognition receptor and a cell adhesion receptor. The neck region is responsible for homo-oligomerization which allows the receptor to bind multivalent ligands with high avidity. A ligand of CD209 is also CD50 (ICAM-3).
References:	*Khoo US, Chan KY, Chan VS, Lin CL: DC-SIGN and L-SIGN: the SIGNs for infection. <i>J Mol Med.</i> 2008 Aug;86(8):861-74. *van Kooyk Y, Geijtenbeek TB: A novel adhesion pathway that regulates dendritic cell trafficking and T cell interactions. <i>Immunol Rev.</i> 2002 Aug;186:47-56. *Geijtenbeek TB, Engering A, Van Kooyk Y: DC-SIGN, a C-type lectin on dendritic cells that unveils many aspects of dendritic cell biology. <i>J Leukoc Biol.</i> 2002 Jun;71(6):921-31. *Ryan EJ, Marshall AJ, Magaletti D, Floyd H, Draves KE, Olson NE, Clark EA: Dendritic cell-associated lectin-1: a novel dendritic cell-associated, C-type lectin-like molecule enhances T cell secretion of IL-4. <i>J Immunol.</i> 2002 Nov 15;169(10):5638-48.

For laboratory research only, not for drug, diagnostic or other use.



Antibodies

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at www.exbio.cz.

For laboratory research only, not for drug, diagnostic or other use.

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic
Tel: +420 261 090 666 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz