



11-643-C025

Monoclonal Antibody to CD361 Purified Antibody (0.025 mg)

Clone:	MEM-216
Isotype:	Mouse IgG1
Specificity:	The mouse monoclonal antibody MEM-216 recognizes CD361 / EVI2B, almost uncharacterized type I transmembrane protein with broad leukocyte expression, mostly in myeloid and B cells. HLDA IX.; WS Code 263
Regulatory Status:	RUO
Immunogen:	Raji cells
Species Reactivity:	Human
Application:	Flow Cytometry Positive control:Raji, Daudi, HL-60 cells, peripheral blood lymphocytes (strongly positive on CD19+ cells) Negative control:Jurkat, U-937 cells Immunoprecipitation
Purity:	> 95% (by SDS-PAGE)
Purification:	Purified by protein-A affinity chromatography
Concentration:	1 mg/ml
Storage Buffer:	Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.4
Storage / Stability:	Store at 2-8°C. Do not freeze. Do not use after expiration date stamped on vial label.
Expiration:	See vial label
Lot Number:	See vial label
Background:	CD361, also known as EVI2B (Ecotropic Viral Integration site 2B) or EVDB, is a poorly characterized type I transmembrane protein, expressed from one of three genes embedded in intron 27b of the neurofibromatosis type 1 (NF1) gene. The DNA strand that is transcribed to produce CD361 is the complementary one to the strand encoding NF1. Murine homolog to human CD361 is associated with ecotropic viral insertions, which have been implicated in the expression of murine myeloid leukemias. CD361 has been also reported to be involved in melanocyte and keratinocyte differentiation. However, it is expressed mainly in peripheral blood and bone marrow.

For laboratory research only, not for drug, diagnostic or other use.



Antibodies

References:

*Viskochil D, Cawthon R, O'Connell P, Xu GF, Stevens J, Culver M, Carey J, White R: The gene encoding the oligodendrocyte-myelin glycoprotein is embedded within the neurofibromatosis type 1 gene. *Mol Cell Biol.* 1991 Feb;11(2):906-12.

*Kaufmann D, Gruener S, Braun F, Stark M, Griesser J, Hoffmeyer S, Bartelt B: EVI2B, a gene lying in an intron of the neurofibromatosis type 1 (NF1) gene, is as the NF1 gene involved in differentiation of melanocytes and keratinocytes and is overexpressed in cells derived from NF1 neurofibromas. *DNA Cell Biol.* 1999 May;18(5):345-56.

*Rudolf-Oliveira RCM, Auat M, Cardoso CC, Santos-Pirath IM, Lange BG, Pires-Silva J, Moraes ACR, Dametto GC, Pirolli MM, Colombo MDHP, Santos-Silva MC: Determination of normal expression patterns of CD86, CD210a, CD261, CD262, CD264, CD358, and CD361 in peripheral blood and bone marrow cells by flow cytometry.

*Mouloungui E, Zver T, Roux C, Amiot C: A protocol to isolate and qualify purified human preantral follicles in cases of acute leukemia, for future clinical applications. *J Ovarian Res.* 2018 Jan 5;11(1):4. doi: 10.1186/s13048-017-0376-6.

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at www.exbio.cz.

For laboratory research only, not for drug, diagnostic or other use.

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic
Tel: +420 261 090 666 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz