



Antibodies

11-647-C025

## Monoclonal Antibody to CD61 Purified Antibody (0.025 mg)

<b>Clone:</b>	VIPL2
<b>Isotype:</b>	Mouse IgG1
<b>Specificity:</b>	The mouse monoclonal antibody VIPL2 recognizes CD61, a 90-110 kDa transmembrane glycoprotein of integrin family, expressed on platelets, megacaryocytes, osteoclasts, endothelial cells and other cell types, including leucocytes and smooth muscle cells. HLDA V.; WS Code 5T-124
<b>Regulatory Status:</b>	RUO
<b>Species Reactivity:</b>	Human, Non-Human Primates
<b>Application:</b>	Flow Cytometry Western Blotting Immunohistochemistry (frozen sections)
<b>Purity:</b>	> 95% (by SDS-PAGE)
<b>Purification:</b>	Purified by protein-A affinity chromatography
<b>Concentration:</b>	1 mg/ml
<b>Storage Buffer:</b>	Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.4
<b>Storage / Stability:</b>	Store at 2-8°C. Do not freeze. Do not use after expiration date stamped on vial label.
<b>Expiration:</b>	See vial label
<b>Lot Number:</b>	See vial label
<b>Background:</b>	CD61 (beta3 integrin) is a transmembrane glycoprotein, which associates with CD41 or CD51 molecules to form heterodimeric adhesion receptors. CD41/CD61 complex is one of the earliest markers of the megakaryocytic lineage. It binds to fibronectin, fibrinogen and von Willebrand factor, and is involved in platelet aggregation. CD51/CD61 complex has similar binding properties and is involved in modulating migration and survival of angiogenic endothelial cells.

**For laboratory research only, not for drug, diagnostic or other use.**



**Antibodies**

- References:**
- \*Dzionek A, Fuchs A, Schmidt P, Cremer S, Zysk M, Miltenyi S, Buck DW, Schmitz J: BDCA-2, BDCA-3, and BDCA-4: three markers for distinct subsets of dendritic cells in human peripheral blood. *J Immunol.* 2000 Dec 1;165(11):6037-46.
  - \*Mondal D, Williams CA, Ali M, Eilers M, Agrawal KC: The HIV-1 Tat protein selectively enhances CXCR4 and inhibits CCR5 expression in megakaryocytic K562 cells. *Exp Biol Med (Maywood).* 2005 Oct;230(9):631-44.
  - \*Williams CA, Mondal D, Agrawal KC: The HIV-1 Tat protein enhances megakaryocytic commitment of K562 cells by facilitating CREB transcription factor coactivation by CBP. *Exp Biol Med (Maywood).* 2005 Dec;230(11):872-84.
  - \*Ciarlet M, Crawford SE, Cheng E, Blutt SE, Rice DA, Bergelson JM, Estes MK: VLA-2 (alpha2beta1) integrin promotes rotavirus entry into cells but is not necessary for rotavirus attachment. *J Virol.* 2002 Feb;76(3):1109-23.
  - \*Barrett L, Dai C, Gamberg J, Gallant M, Grant M: Circulating CD14-CD36+ peripheral blood mononuclear cells constitutively produce interleukin-10. *J Leukoc Biol.* 2007 Jul;82(1):152-60.
  - \*Roberts MS, Woods AJ, Dale TC, Van Der Sluijs P, Norman JC: Protein kinase B/Akt acts via glycogen synthase kinase 3 to regulate recycling of alpha v beta 3 and alpha 5 beta 1 integrins. *Mol Cell Biol.* 2004 Feb;24(4):1505-15.

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at [www.exbio.cz](http://www.exbio.cz).

**For laboratory research only, not for drug, diagnostic or other use.**

---

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic  
Tel: +420 261 090 666 | Fax: +420 261 090 660 | [orders@exbio.cz](mailto:orders@exbio.cz) | [www.exbio.cz](http://www.exbio.cz)