



11-650-C025

## Monoclonal Antibody to CD314 Purified Antibody (0.025 mg)

<b>Clone:</b>	1D11
<b>Isotype:</b>	Mouse IgG1
<b>Specificity:</b>	The mouse monoclonal antibody 1D11 recognizes CD314 / NKG2D, a 42 kDa C-type lectin-like activating receptor expressed by NK cells, gamma/delta T cells, and CD8+ T cells.
<b>Regulatory Status:</b>	RUO
<b>Immunogen:</b>	NKL cell line
<b>Species Reactivity:</b>	Human
<b>Application:</b>	Flow Cytometry Immunoprecipitation Immunohistochemistry (frozen sections) Functional Application blocking of ligand binding
<b>Purity:</b>	> 95% (by SDS-PAGE)
<b>Purification:</b>	Purified by protein-A affinity chromatography
<b>Concentration:</b>	1 mg/ml
<b>Storage Buffer:</b>	Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.4
<b>Storage / Stability:</b>	Store at 2-8°C. Do not freeze. Do not use after expiration date stamped on vial label.
<b>Expiration:</b>	See vial label
<b>Lot Number:</b>	See vial label
<b>Background:</b>	CD314, also known as NKG2D (natural killer receptor G2D) or KLRK1 (killer cell lectin-like receptor subfamily K, member 1), is a homodimeric C-type lectin-like activating receptor and costimulator with type II membrane orientation (C terminus extracellular). CD314 homodimers are associated with DAP10, a membrane adaptor protein that signals similar to CD28 by recruitment of phosphatidylinositol 3-kinase. Engagement of CD314 amplifies antigen-specific T cell responses in CD314-positive T cell populations. In NK cells, CD314 is a primary activating receptor. As CD314 ligands the MHC class-I chain-related proteins A and B (MICA, MICB) and UL16-binding proteins (ULBPs) have been identified.

**For laboratory research only, not for drug, diagnostic or other use.**

**Antibodies****References:**

- \*Bauer S, Groh V, Wu J, Steinle A, Phillips JH, Lanier LL, Spies T: Activation of NK cells and T cells by NKG2D, a receptor for stress-inducible MICA. *Science*. 1999 Jul 30;285(5428):727-9.
- \*Wu J, Cherwinski H, Spies T, Phillips JH, Lanier LL: DAP10 and DAP12 form distinct, but functionally cooperative, receptor complexes in natural killer cells. *J Exp Med*. 2000 Oct 2;192(7):1059-68.
- \*Wu J, Groh V, Spies T: T cell antigen receptor engagement and specificity in the recognition of stress-inducible MHC class I-related chains by human epithelial gamma delta T cells. *J Immunol*. 2002 Aug 1;169(3):1236-40.
- \*Sangiolo D, Martinuzzi E, Todorovic M, Vitaggio K, Vallario A, Jordaney N, Carnevale-Schianca F, Capaldi A, Geuna M, Casorzo L, Nash RA, Aglietta M, Cignetti A: Alloreactivity and anti-tumor activity segregate within two distinct subsets of cytokine-induced killer (CIK) cells: implications for their infusion across major HLA barriers. *Int Immunol*. 2008 Jul;20(7):841-8.
- \*Hasenkamp J, Borgerding A, Uhrberg M, Falk C, Chapuy B, Wulf G, Jung W, Trümper L, Glass B: Self-tolerance of human natural killer cells lacking self-HLA-specific inhibitory receptors. *Scand J Immunol*. 2008 Mar;67(3):218-29.
- \*Ebert LM, Meuter S, Moser B: Homing and function of human skin gammadelta T cells and NK cells: relevance for tumor surveillance. *J Immunol*. 2006 Apr 1;176(7):4331-6.
- \*Valencia J, Hernández-López C, Martínez VG, Hidalgo L, Zapata AG, Vicente A, Varas A, Sacedón R: Transient beta-catenin stabilization modifies lineage output from human thymic CD34+CD1a- progenitors. *J Leukoc Biol*. 2010 Mar;87(3):405-14.

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at [www.exbio.cz](http://www.exbio.cz).

**For laboratory research only, not for drug, diagnostic or other use.**

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic  
Tel: +420 261 090 666 | Fax: +420 261 090 660 | [orders@exbio.cz](mailto:orders@exbio.cz) | [www.exbio.cz](http://www.exbio.cz)