



Antibodies

11-667-C100

Monoclonal Antibody to CD62P Purified Antibody (0.1 mg)

Clone:	AK4
Isotype:	Mouse IgG1
Specificity:	The antibody AK4 recognizes CD62P (P-selectin), a 140 kD single chain type I transmembrane glycoprotein present in secretory alpha-granules in platelets, in Weibel-Palade bodies in endothelial cells and in megakaryocytes; it is relocated to the plasma membrane upon activation. HLDA VI; WS Code P-44
Regulatory Status:	RUO
Immunogen:	Human platelets
Species Reactivity:	Human, Non-Human Primates
Application:	Flow Cytometry Recommended dilution: 1.5 µg/ml Immunoprecipitation Western Blotting Immunocytochemistry Functional Application Blocking
Purity:	> 95% (by SDS-PAGE)
Purification:	Purified by protein-A affinity chromatography
Concentration:	1 mg/ml
Storage Buffer:	Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.4
Storage / Stability:	Store at 2-8°C. Do not freeze. Do not use after expiration date stamped on vial label.
Expiration:	See vial label
Lot Number:	See vial label
Background:	CD62P (P-selectin) is an adhesion glycoprotein that is expressed on platelets and endothelial cells upon their activation. Interaction between CD62P and its mucin-like ligand PSGL-1 (P-selectin glycoprotein ligand-1) expressed on the microvilli of most leukocytes supports leukocyte rolling along postkapillary venules at the earliest time of inflammation. Both CD62P and PSGL-1 are extended glycoproteins that form homodimers. CD62P dimerization is probably mediated through interactions of the transmembrane domains and stabilizes leukocyte tethering and rolling, probably by increasing rebinding within a bond cluster.

For laboratory research only, not for drug, diagnostic or other use.



Antibodies

- References:**
- *Skinner MP, Lucas CM, Burns GF, Chesterman CN, Berndt MC: GMP-140 binding to neutrophils is inhibited by sulfated glycans. *J Biol Chem.* 1991 Mar 25;266(9):5371-4.
 - *Dunlop LC, Skinner MP, Bendall LJ, Favalaro EJ, Castaldi PA, Gorman JJ, Gamble JR, Vadas MA, Berndt MC: Characterization of GMP-140 (P-selectin) as a circulating plasma protein. *J Exp Med.* 1992 Apr 1;175(4):1147-50.
 - *Holme PA, Müller F, Solum NO, Brosstad F, Frøland SS, Aukrust P: Enhanced activation of platelets with abnormal release of RANTES in human immunodeficiency virus type 1 infection. *FASEB J.* 1998 Jan;12(1):79-89.
 - *Kowalska MA, Ratajczak J, Hoxie J, Brass LF, Gewirtz A, Poncz M, Ratajczak MZ: Megakaryocyte precursors, megakaryocytes and platelets express the HIV co-receptor CXCR4 on their surface: determination of response to stromal-derived factor-1 by megakaryocytes and platelets. *Br J Haematol.* 1999 Feb;104(2):220-9.
 - *Ludwig RJ, Schultz JE, Boehncke WH, Podda M, Tandi C, Krombach F, Baatz H, Kaufmann R, von Andrian UH, Zollner TM: Activated, not resting, platelets increase leukocyte rolling in murine skin utilizing a distinct set of adhesion molecules. *J Invest Dermatol.* 2004 Mar;122(3):830-6.

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at www.exbio.cz.

For laboratory research only, not for drug, diagnostic or other use.

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic
Tel: +420 261 090 666 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz