



Antibodies

11-669-C100

Monoclonal Antibody to CD205 Purified Antibody (0.1 mg)

Clone:	HD30
Isotype:	Mouse IgG1
Specificity:	The mouse monoclonal antibody HD30 recognizes CD205, an approx. 200 kDa C-type lectin transmembrane protein of the MMR (macrophage mannose receptor) family, expressed at high levels on dendritic cells and thymic epithelial cells, and at low levels on lymphocytes, NK cells and monocytes.
Regulatory Status:	RUO
Immunogen:	Recombinant Fc-tagged human CD205
Species Reactivity:	Human
Application:	Flow Cytometry Western Blotting Application note: only non-denatured samples Immunohistochemistry (frozen sections)
Purity:	> 95% (by SDS-PAGE)
Purification:	Purified by protein-A affinity chromatography
Concentration:	1 mg/ml
Storage Buffer:	Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.4
Storage / Stability:	Store at 2-8°C. Do not freeze. Do not use after expiration date stamped on vial label.
Expiration:	See vial label
Lot Number:	See vial label
Background:	CD205, also known as DEC-205, is an endocytic receptor of macrophage mannose receptor family. This 205 kDa C-type lectin transmembrane protein mediates adsorptive uptake and its intracellular domain contains coated pit localization sequence and distal acidic motif, which is required for recycling beyond early endosomes through deeper MHC II+ late endosomes and lysosomes. This unique pathway of receptor-mediated uptake proves to be necessary for presentation of antigenic peptides at low doses of ligand. CD205 is responsible for uptake and processing of captured antigens for dendritic cells.

For laboratory research only, not for drug, diagnostic or other use.



Antibodies

- References:**
- *Gurer C, Strowig T, Brilot F, Pack M, Trumfheller C, Arrey F, Park CG, Steinman RM, Münz C: Targeting the nuclear antigen 1 of Epstein-Barr virus to the human endocytic receptor DEC-205 stimulates protective T-cell responses. *Blood*. 2008 Aug 15;112(4):1231-9.
 - *Guo M, Gong S, Maric S, Misulovin Z, Pack M, Mahnke K, Nussenzweig MC, Steinman RM: A monoclonal antibody to the DEC-205 endocytosis receptor on human dendritic cells. *Hum Immunol*. 2000 Aug;61(8):729-38.
 - *Mahnke K, Guo M, Lee S, Sepulveda H, Swain SL, Nussenzweig M, Steinman RM: The dendritic cell receptor for endocytosis, DEC-205, can recycle and enhance antigen presentation via major histocompatibility complex class II-positive lysosomal compartments. *J Cell Biol*. 2000 Oct 30;151(3):673-84.
 - *Park CG, Rodriguez A, Ueta H, Lee H, Pack M, Matsuno K, Steinman RM: Generation of anti-human DEC205/CD205 monoclonal antibodies that recognize epitopes conserved in different mammals. 2012 Mar 30;377(1-2):15-22.

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at www.exbio.cz.

For laboratory research only, not for drug, diagnostic or other use.

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic
Tel: +420 261 090 666 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz