



11-697-C100

Monoclonal Antibody to CD148 Purified Antibody (0.1 mg)

Clone:	MEM-CD148/05
Isotype:	Mouse IgG2b
Specificity:	The mouse monoclonal antibody MEM-CD148/05 recognizes CD148, a highly glycosylated up to 250 kDa receptor-like protein tyrosin phosphatase expressed mainly in lymphocytes, myeloid cells and epithelial cells.
Regulatory Status:	RUO
Immunogen:	Human recombinant CD148 (amino acids 1-444)
Species Reactivity:	Human
Application:	Flow Cytometry
Purity:	> 95% (by SDS-PAGE)
Purification:	Purified by protein-A affinity chromatography
Concentration:	1 mg/ml
Storage Buffer:	Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.4
Storage / Stability:	Store at 2-8°C. Do not freeze. Do not use after expiration date stamped on vial label.
Expiration:	See vial label
Lot Number:	See vial label
Background:	CD148 (also known as HPTP-eta or DEP-1) is a transmembrane protein tyrosin phosphatase containing eight fibronectin type III extracellular domains. This protein is known to inhibit transduction of mitogenic signals in non-hematopoietic cells (fibroblasts, epithelial cells), and signal transduction downstream of T cell receptor, however, it also augments immunoreceptor signaling in B cells and macrophages via dephosphorylating C-terminal tyrosine of Src-family tyrosine kinases. CD148 expression increases after in vitro activation of peripheral blood leucocytes. It can be also used as marker of the most mature human thymocytes, and leukemic cells corresponding to this stadium of thymocyte differentiation. In contrast, in mice the CD148 expression sharply drops through the double positive stage to the single positive thymocytes.
References:	*Stepanek O, Kalina T, Draber P, Skopcova T, Svojgr K, Angelisova P, Horejsi V, Weiss A, Brdicka T: Regulation of Src family kinases involved in T cell receptor signaling by protein-tyrosine phosphatase CD148. J Biol Chem. 2011 Jun 24;286(25):22101-12.

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at www.exbio.cz.

For laboratory research only, not for drug, diagnostic or other use.

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic
Tel: +420 261 090 666 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz