

11-728-C100

Monoclonal Antibody to CD300e Purified Antibody (0.1 mg)

Clone: UP-H2

Isotype: Mouse IgG1

Specificity: The mouse monoclonal antibody UP-H2 recognizes CD300e / IREM-2, a 32 kDa

glycoprotein expressed by mature monocytes and peripheral blood myeloid

dendritic cells.

Regulatory Status: RUO

Immunogen: CD300e-HA-transfected cells

Species Reactivity: Human

Application: Flow Cytometry

Immunoprecipitation Western Blotting Functional Application

stimulation

Purity: > 95% (by SDS-PAGE)

Purification: Purified by protein-A affinity chromatography

Concentration: 1 mg/ml

Storage Buffer: Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.4

Storage / Stability: Store at 2-8°C. Do not freeze. Do not use after expiration date stamped on vial

label.

Expiration: See vial label

Lot Number: See vial label

Background: CD300e / IREM-2 (immune receptor expressed by myeloid cells 2), also known as

CLM2 or LMIR6, is a monomeric transmembrane glycoprotein with a single extracellular immunoglobulin-like domain. Intracellularly it associates with DAP-12, an ITAM-containing adaptor molecule. CD300e is expressed on mature monocytes and peripheral blood myeloid dendritic cells. Its crosslinking leads to release of

pro-inflammatory cytokines, and increased expression of activation markers.

References: *Clark GJ, Ju X, Tate C, Hart DN: The CD300 family of molecules are

evolutionarily significant regulators of leukocyte functions. Trends Immunol. 2009

May;30(5):209-17.

*Aguilar H, Alvarez-Errico D, García-Montero AC, Orfao A, Sayós J, López-Botet M: Molecular characterization of a novel immune receptor restricted to the

monocytic lineage. J Immunol. 2004 Dec 1;173(11):6703-11.

*Brckalo T, Calzetti F, Pérez-Cabezas B, Borràs FE, Cassatella MA, López-Botet M: Functional analysis of the CD300e receptor in human monocytes

and myeloid dendritic cells. Eur J Immunol. 2010 Mar;40(3):722-32.

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at www.exbio.cz.

For laboratory research only, not for drug, diagnostic or other use.