



11-729-C100

Monoclonal Antibody to CD161 Purified Antibody (0.1 mg)

Clone:	HP-3G10
Isotype:	Mouse IgG1
Specificity:	The mouse monoclonal antibody HP-3G10 recognizes CD161, a type II transmembrane C-type lectin receptor, expressed on the plasma membrane of NK cells, dendritic cells, activated monocytes and a subset of T cells as a disulphide-linked homodimer.
Regulatory Status:	RUO
Immunogen:	human NK cells
Species Reactivity:	Human, Non-Human Primates
Application:	Flow Cytometry Western Blotting
Purity:	> 95% (by SDS-PAGE)
Purification:	Purified by protein-A affinity chromatography
Concentration:	1 mg/ml
Storage Buffer:	Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.4
Storage / Stability:	Store at 2-8°C. Do not freeze. Do not use after expiration date stamped on vial label.
Expiration:	See vial label
Lot Number:	See vial label
Background:	CD161, also known as Nkrp1 (natural killer receptor protein 1) or Klrk1 (killer cell lectin-like receptor subfamily b member 1), is a disulphide-linked homodimeric receptor, which is involved in regulation of NK cell and NKT cell function. It is expressed on a majority of NK cells, NKT cells, and e.g. Th17 cells and CD3+ thymocytes. Although rat CD161 has three isoforms (a, b, c), the human CD161 is expressed as one isoform.

For laboratory research only, not for drug, diagnostic or other use.

**Antibodies**

- References:**
- *Poza D, Valés-Gómez M, Mavaddat N, Williamson SC, Chisholm SE, Reyburn H: CD161 (human NKR-P1A) signaling in NK cells involves the activation of acid sphingomyelinase. *J Immunol.* 2006 Feb 15;176(4):2397-406.
 - *Germain C, Meier A, Jensen T, Knapnougel P, Poupon G, Lazzari A, Neisig A, Håkansson K, Dong T, Wagtmann N, Galsgaard ED, Spee P, Braud VM: Induction of lectin-like transcript 1 (LLT1) protein cell surface expression by pathogens and interferon- γ ; contributes to modulate immune responses. *J Biol Chem.* 2011 Nov 4;286(44):37964-75
 - *Wong EB, Akilimali NA, Govender P, Sullivan ZA, Cosgrove C, Pillay M, Lewensohn DM, Bishai WR, Walker BD, Ndung'u T, Klenerman P, Kasproicz VO: Low Levels of Peripheral CD161⁺⁺CD8⁺ Mucosal Associated Invariant T (MAIT) Cells Are Found in HIV and HIV/TB Co-Infection. *PLoS One.* 2013 Dec 31;8(12):e83474
 - *Birgit Fogal, Tai Yi, Chen Wang, Deepak A. Rao, Amir Lebastchi, Sanjay Kulkarni, George Tellides, Jordan S. Pober: Neutralizing IL-6 reduces human arterial allograft rejection by allowing emergence of CD161(+) CD4(+) T regulatory cells. *J Immunol.* 2011 December 15; 187(12): 6268–6280.
 - *Tresoldi E, Dell'Albani I, Stabilini A, Jofra T, Valle A, Gagliani N, Bondanza A, Roncarolo MG, Battaglia M: Stability of human rapamycin-expanded CD4⁺CD25⁺ T regulatory cells. *Haematologica.* 2011 Sep;96(9):1357-65.
 - *Goetzl EJ, Huang MC, Kon J, Patel K, Schwartz JB, Fast K, Ferrucci L, Madara K, Taub DD, Longo DL: Gender specificity of altered human immune cytokine profiles in aging. *FASEB J.* 2010 Sep;24(9):3580-9.

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at www.exbio.cz.

For laboratory research only, not for drug, diagnostic or other use.

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic
Tel: +420 261 090 666 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz