



11-732-C100

Monoclonal Antibody to CD42a Purified Antibody (0.1 mg)

Clone:	GR-P
Isotype:	Mouse IgG1
Specificity:	The mouse monoclonal antibody GR-P (also known as GRP-P) recognizes CD42a (glycoprotein 9), a 22 kDa transmembrane protein constitutively expressed on megakaryocytes and platelets. HLDA IV.; WS Code P 35
Regulatory Status:	RUO
Immunogen:	Human acute lymphoblastic leukemia cells
Species Reactivity:	Human, Canine (Dog)
Application:	Flow Cytometry
Purity:	> 95% (by SDS-PAGE)
Purification:	Purified by protein-A affinity chromatography
Concentration:	1 mg/ml
Storage Buffer:	Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.4
Storage / Stability:	Store at 2-8°C. Do not freeze. Do not use after expiration date stamped on vial label.
Expiration:	See vial label
Lot Number:	See vial label
Background:	CD42a, also known as Glycoprotein 9 (GPIX), composes together with GPIb alpha, GPIb beta and GPV the GPIb-IX-V receptor complex critical in the process of platelet-rich thrombus formation by tethering the platelet to a thrombogenic surface. CD42b binds to von Willebrand factor (VWF) exposed at a site of vascular injury, as well as to thrombin, coagulation factors XI and XII, high molecular weight kininogen, TSP-1, integrin Mac-1 and P-selectin. Defects in the gene encoding CD42a are a cause of Bernard-Soulier syndrome, also known as giant platelet disease. These patients have unusually large platelets and have a clinical bleeding tendency.

For laboratory research only, not for drug, diagnostic or other use.



Antibodies

- References:**
- *Leukocyte Typing IV., Knapp W. et al. (Eds.), Oxford University Press (1989)
 - *Harding SA, Din JN, Sarma J, Josephs DH, Fox KA, Newby DE: Promotion of proinflammatory interactions between platelets and monocytes by unfractionated heparin. *Heart*. 2006 Nov;92(11):1635-8.
 - *Brown SB, Clarke MC, Magowan L, Sanderson H, Savill J: Constitutive death of platelets leading to scavenger receptor-mediated phagocytosis. A caspase-independent cell clearance program. *J Biol Chem*. 2000 Feb 25;275(8):5987-96.
 - *Vettore S, Scandellari R, Moro S, Lombardi AM, Scapin M, Randi ML, Fabris F: Novel point mutation in a leucine-rich repeat of the GPIIb/IIIa chain of the platelet von Willebrand factor receptor, GPIIb/IX/V, resulting in an inherited dominant form of Bernard-Soulier syndrome affecting two unrelated families: the N41H variant. *Haematologica*. 2008 Nov;93(11):1743-7.
 - *Din JN, Aftab SM, Jubb AW, Carnegy FH, Lyall K, Sarma J, Newby DE, Flapan AD: Effect of moderate walnut consumption on lipid profile, arterial stiffness and platelet activation in humans. *Eur J Clin Nutr*. 2011 Feb;65(2):234-9.

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at www.exbio.cz.

For laboratory research only, not for drug, diagnostic or other use.

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic
Tel: +420 261 090 666 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz