



11-735-C025

Monoclonal Antibody to CD197 / CCR7 (mouse) Purified Antibody (0.025 mg)

Clone:	4B12
Isotype:	Rat IgG2a
Specificity:	The rat monoclonal antibody 4B12 recognizes murine CD197 / CCR7 (chemokine receptor 7), a 43 kDa G-protein-coupled receptor for chemokines CCL19 and CCL21.
Regulatory Status:	RUO
Immunogen:	Murine CD197-transfected RBL-2H3 cells
Species Reactivity:	Mouse
Application:	Flow Cytometry Immunoprecipitation
Purity:	> 95% (by SDS-PAGE)
Purification:	Purified by protein-G affinity chromatography
Concentration:	1 mg/ml
Storage Buffer:	Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.4
Storage / Stability:	Store at 2-8°C. Do not freeze. Do not use after expiration date stamped on vial label.
Expiration:	See vial label
Lot Number:	See vial label
Background:	CD197 / CCR7 (chemokine receptor 7) is a seven membrane-spanning protein serving as a receptor for chemokines CCL19 and CCL21. It is expressed on most naive T cells, some hematopoietic stem cells, mature dendritic cells, NK cells and some memory T cells and B cells subsets. CD197 plays important roles in development of thymocytes, and regulating the recirculation and homing of lymphocytes and dendritic cells to secondary lymphoid organs.
References:	*Singh V, Gowthaman U, Jain S, Parihar P, Banskar S, Gupta P, Gupta UD, Agrewala JN: Coadministration of interleukins 7 and 15 with bacille Calmette-Guérin mounts enduring T cell memory response against Mycobacterium tuberculosis. <i>J Infect Dis.</i> 2010 Aug 15;202(3):480-9 *Adachi K, Osada Y, Nakamura R, Tamada K, Hamano S: Unique T Cells with Unconventional Cytokine Profiles Induced in the Livers of Mice during Schistosoma mansoni Infection. <i>PLoS One.</i> 2013 Dec 16;8(12):e82698. *Huggins A, Paschalidis N, Flower RJ, Perretti M, D'Acquisto F: Annexin-1-deficient dendritic cells acquire a mature phenotype during differentiation. <i>FASEB J.</i> 2009 Apr;23(4):985-96 *Eisenbarth SC, Williams A, Colegio OR, Meng H, Strowig T, Rongvaux A, Henao-Mejia J, Thaiss CA, Joly S, Gonzalez DG, Xu L, Zenewicz LA, Haberman AM, Elinav E, Kleinstein SH, Sutterwala FS, Flavell RA: NLRP10 is a NOD-like receptor essential to initiate adaptive immunity by dendritic cells. <i>Nature.</i> 2012 Apr 25;484(7395):510-3

For laboratory research only, not for drug, diagnostic or other use.



Antibodies

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at www.exbio.cz.

For laboratory research only, not for drug, diagnostic or other use.

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic
Tel: +420 261 090 666 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz