



Antibodies

11-740-C025

Monoclonal Antibody to CD85j / ILT2 Purified Antibody (0.025 mg)

Clone:	GHI/75
Isotype:	Mouse IgG2b
Specificity:	The mouse monoclonal antibody GHI/75 recognizes CD85j / ILT2, an 110-120 kDa membrane glycoprotein expressed strongly on plasma cells, moderately on circulating B cells, and weakly on monocytes. It is also expressed on T cell and NK cell subsets (variable, individual).
Regulatory Status:	RUO
Immunogen:	Hairy cell leukaemia cells
Species Reactivity:	Human
Application:	Flow Cytometry Immunoprecipitation Western Blotting Functional Application blocking HLA-G induced TGF-beta1 production
Purity:	> 95% (by SDS-PAGE)
Purification:	Purified by protein-A affinity chromatography
Concentration:	1 mg/ml
Storage Buffer:	Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.4
Storage / Stability:	Store at 2-8°C. Do not freeze. Do not use after expiration date stamped on vial label.
Expiration:	See vial label
Lot Number:	See vial label
Background:	CD85j, also known as ILT-2 (Ig-like transcript 2), LIR-1 (leukocyte Ig-like receptor 1), or LILRB1 (leukocyte Ig-like receptor B1), is a member of Ig superfamily transmembrane glycoproteins named CD85. The CD85j protein is expressed on several types of immune cells (plasma cells, B cells, monocytes, T and NK cell subsets) where it binds to MHC class I molecules on antigen-presenting cells and transduces a negative signal that inhibits stimulation of an immune response. It is thought to control inflammatory responses and cytotoxicity to help focus the immune response and limit autoreactivity.

For laboratory research only, not for drug, diagnostic or other use.



Antibodies

- References:**
- *Pulford K, Micklem K, Thomas J, Jones M, Mason DY: A 72-kD B cell-associated surface glycoprotein expressed at high levels in hairy cell leukaemia and plasma cell neoplasms. *Clin Exp Immunol.* 1991 Sep;85(3):429-35.
 - *Banham AH, Colonna M, Cella M, Micklem KJ, Pulford K, Willis AC, Mason DY: Identification of the CD85 antigen as ILT2, an inhibitory MHC class I receptor of the immunoglobulin superfamily. *J Leukoc Biol.* 1999 Jun;65(6):841-5.
 - *Leukocyte Typing V., Schlossman S. et al. (Eds.), Oxford University Press (1995).
 - *Lo Monaco E, Tremante E, Cerboni C, Melucci E, Sibilio L, Zingoni A, Nicotra MR, Natali PG, Giacomini P: Human leukocyte antigen E contributes to protect tumor cells from lysis by natural killer cells. *Neoplasia.* 2011 Sep;13(9):822-30.
 - *Riteau B, Menier C, Khalil-Daher I, Martinozzi S, Pla M, Dausset J, Carosella ED, Rouas-Freiss N: HLA-G1 co-expression boosts the HLA class I-mediated NK lysis inhibition. *Int Immunol.* 2001 Feb;13(2):193-201.

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at www.exbio.cz.

For laboratory research only, not for drug, diagnostic or other use.