



11-751-C025

Monoclonal Antibody to CD32 Purified Antibody (0.025 mg)

Clone:	3D3
Isotype:	Mouse IgG1
Specificity:	The mouse monoclonal antibody 3D3 recognizes CD32, a 40 kDa polymorphic transmembrane glycoprotein serving as the low affinity receptor for aggregated IgG. This antibody recognizes CD32 on B cells of all donors, but on platelets, monocytes, and granulocytes of only some donors (131R variant, but not 131H variant).
Regulatory Status:	RUO
Immunogen:	purified glycosylated recombinant human FcγRIIIa2
Species Reactivity:	Human
Application:	Flow Cytometry Mass Cytometry
Purity:	> 95% (by SDS-PAGE)
Purification:	Purified by protein-A affinity chromatography
Concentration:	1 mg/ml
Storage Buffer:	Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.4
Storage / Stability:	Store at 2-8°C. Do not freeze. Do not use after expiration date stamped on vial label.
Expiration:	See vial label
Lot Number:	See vial label
Background:	CD32 (FcγRIII) is a low affinity receptor for aggregated IgG. It is strongly expressed on monocytes, granulocytes, myeloid and myeloblastic cell lines, and weakly on B cells, CD34+ bone marrow cells, and resting and activated platelets. After binding its ligand, CD32 induces IgG-mediated phagocytosis and oxidative burst in monocytes and neutrophils, whereas in B cells it mediates a negative signal. This polymorphic transmembrane glycoprotein is expressed not only in the activating (CD32a) and inhibitory isoform (CD32b), but also in individual variants with differing avidities for IgG subtypes (e.g. the CD32a131R and CD32a131H allotypes).

For laboratory research only, not for drug, diagnostic or other use.

**Antibodies****References:**

*Vely F, Gruel N, Moncuit J, Cochet O, Rouard H, Dare S, Galon J, Sautes C, Fridman WH, Teillaud JL: A new set of monoclonal antibodies against human Fc gamma RII (CD32) and Fc gamma RIII (CD16): characterization and use in various assays. *Hybridoma*. 1997 Dec;16(6):519-28.

*Boruchov AM, Heller G, Veri MC, Bonvini E, Ravetch JV, Young JW: Activating and inhibitory IgG Fc receptors on human DCs mediate opposing functions. *J Clin Invest*. 2005 Oct;115(10):2914-23.

*Dutertre CA, Bonnin-Gélizé E, Pulford K, Bourel D, Fridman WH, Teillaud JL: A novel subset of NK cells expressing high levels of inhibitory Fc gamma RIIb modulating antibody-dependent function. *J Leukoc Biol*. 2008 Dec;84(6):1511-20

*Os A, Bürgler S, Ribes AP, Funderud A, Wang D, Thompson KM, Tj#248;nnfjord GE, Bogen B, Munthe LA: Chronic lymphocytic leukemia cells are activated and proliferate in response to specific T helper cells. *Cell Rep*. 2013 Aug 15;4(3):566-77.

*Dai X, Jayapal M, Tay HK, Reghunathan R, Lin G, Too CT, Lim YT, Chan SH, Kemeny DM, Floto RA, Smith KG, Melendez AJ, MacAry PA: Differential signal transduction, membrane trafficking, and immune effector functions mediated by Fc gamma RI versus Fc gamma RIIa. *Blood*. 2009 Jul 9;114(2):318-27.

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at www.exbio.cz.

For laboratory research only, not for drug, diagnostic or other use.

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic
Tel: +420 261 090 666 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz