



11-753-C025

## Monoclonal Antibody to CD95 / Fas Purified Antibody (0.025 mg)

<b>Clone:</b>	EOS9.1
<b>Isotype:</b>	Mouse IgM
<b>Specificity:</b>	The mouse monoclonal antibody EOS9.1 recognizes CD95 (Fas/APO-1), a 46 kDa glycoprotein of the tumour necrosis factor/nerve growth factor (TNF/NGF) receptor superfamily, expressed on a variety of normal and neoplastic cells.
<b>Regulatory Status:</b>	RUO
<b>Immunogen:</b>	P815 cells transfected with human CD95
<b>Species Reactivity:</b>	Human
<b>Application:</b>	Functional Application in vitro induction of apoptosis
<b>Purity:</b>	> 95% (by SDS-PAGE)
<b>Purification:</b>	Purified by precipitation and chromatography
<b>Concentration:</b>	1 mg/ml
<b>Storage Buffer:</b>	Tris buffered saline (TBS) with 15 mM sodium azide, approx. pH 8.0
<b>Storage / Stability:</b>	Store at 2-8°C. Do not freeze. Do not use after expiration date stamped on vial label.
<b>Expiration:</b>	See vial label
<b>Lot Number:</b>	See vial label
<b>Background:</b>	CD95 (Fas, APO-1), a 46 kDa transmembrane glycoprotein, is a cell death receptor of the TNFR superfamily. Stimulation of CD95 results in aggregation of its intracellular death domains, formation of the death-inducing signaling complex (DISC) and activation of caspases. In type I cells caspase 3 is activated by high amounts of caspase 8 generated at the DISC, in type II cells low concentration of caspase 8 activates pathway leading to the release of cytochrome c from mitochondria and activation of caspase 3 by cytochrome c. Besides its roles in induction of apoptosis, Fas also triggers pro-inflammatory cytokine responses.
<b>References:</b>	*Conejo-Garcia JR, Benencia F, Courreges MC, Gimotty PA, Khang E, Buckanovich RJ, Frauwirth KA, Zhang L, Katsaros D, Thompson CB, Levine B, Coukos G: Ovarian carcinoma expresses the NKG2D ligand Letal and promotes the survival and expansion of CD28- antitumor T cells. <i>Cancer Res.</i> 2004 Mar 15;64(6):2175-82. *Kasper HU, Konze E, Kern M, Stippel DL: CD95 and TNF-induced apoptosis in liver metastases of colorectal carcinoma. <i>In Vivo.</i> 2010 Sep-Oct;24(5):653-7. *Matsuoka K, Kim HT, McDonough S, Bascug G, Warshauer B, Koreth J, Cutler C, Ho VT, Alyea EP, Antin JH, Soiffer RJ, Ritz J: Altered regulatory T cell homeostasis in patients with CD4+ lymphopenia following allogeneic hematopoietic stem cell transplantation. <i>J Clin Invest.</i> 2010 May;120(5):1479-93 *Desbarats J, Birge RB, Mimouni-Rongy M, Weinstein DE, Palerme JS, Newell MK: Fas engagement induces neurite growth through ERK activation and p35 upregulation. <i>Nat Cell Biol.</i> 2003 Feb;5(2):118-25.

**For laboratory research only, not for drug, diagnostic or other use.**



**Antibodies**

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at [www.exbio.cz](http://www.exbio.cz).

**For laboratory research only, not for drug, diagnostic or other use.**

---

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic  
Tel: +420 261 090 666 | Fax: +420 261 090 660 | [orders@exbio.cz](mailto:orders@exbio.cz) | [www.exbio.cz](http://www.exbio.cz)