



11-761-C100

## Monoclonal Antibody to CD1b Purified Antibody (0.1 mg)

<b>Clone:</b>	SN13
<b>Isotype:</b>	Mouse IgG1
<b>Specificity:</b>	The mouse monoclonal antibody SN13 (also known as K5-1B8) recognizes CD1b, a 44 kDa type I glycoprotein associated with beta2-microglobulin. It is expressed on dendritic cells, Langerhans cells, thymocytes, and T acute lymphoblastic leukemia cells.
<b>Regulatory Status:</b>	RUO
<b>Immunogen:</b>	A cell membrane antigen preparation that was isolated from normal human thymocytes
<b>Species Reactivity:</b>	Human
<b>Application:</b>	Flow Cytometry Immunoprecipitation Immunohistochemistry (paraffin sections) Immunohistochemistry (frozen sections)
<b>Purity:</b>	> 95% (by SDS-PAGE)
<b>Purification:</b>	Purified by protein-A affinity chromatography
<b>Concentration:</b>	1 mg/ml
<b>Storage Buffer:</b>	Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.4
<b>Storage / Stability:</b>	Store at 2-8°C. Do not freeze. Do not use after expiration date stamped on vial label.
<b>Expiration:</b>	See vial label
<b>Lot Number:</b>	See vial label
<b>Background:</b>	CD1b (also known as R1) together with CD1a and c, belongs to group 1 of CD1 antigens. These non-classical MHC-like glycoproteins serve as antigen-presenting molecules for a subset of T cells that responds to specific lipids and glycolipids found in the cell walls of bacterial pathogens or self-glycolipid antigens such as gangliosides, and they have also roles in antiviral immunity. The trafficking routes of the particular CD1 types differ and correspond to their ability to bind and present different groups of antigens. Besides non-peptide glycolipid antigen presentation to CD1-restricted T cells, CD1b has been implicated in thymocyte development.

**For laboratory research only, not for drug, diagnostic or other use.**



**Antibodies**

**References:**

- \*Leukocyte Typing VI., Kishimoto T. et al. (Eds.), Garland Publishing Inc. (1997).
- \*Giuliani A, Prete SP, Graziani G, Aquino A, Balduzzi A, Sugita M, Brenner MB, Iona E, Fattorini L, Orefici G, Porcelli SA, Bonmassar E: Influence of Mycobacterium bovis bacillus Calmette Guérin on in vitro induction of CD1 molecules in human adherent mononuclear cells. Infect Immun. 2001 Dec;69(12):7461-70.
- \*Tentori L, Graziani G, Porcelli SA, Sugita M, Brenner MB, Madaio R, Bonmassar E, Giuliani A, Aquino A: Rifampin increases cytokine-induced expression of the CD1b molecule in human peripheral blood monocytes. Antimicrob Agents Chemother. 1998 Mar;42(3):550-4.
- \*Hayes SM, Knight KL: Group 1 CD1 genes in rabbit. J Immunol. 2001 Jan 1;166(1):403-10.

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at [www.exbio.cz](http://www.exbio.cz).

**For laboratory research only, not for drug, diagnostic or other use.**

---

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic  
Tel: +420 261 090 666 | Fax: +420 261 090 660 | [orders@exbio.cz](mailto:orders@exbio.cz) | [www.exbio.cz](http://www.exbio.cz)