



11-778-C025

Monoclonal Antibody to CD101 Purified Antibody (0.025 mg)

Clone:	BB27
Isotype:	Mouse IgG1
Specificity:	The mouse monoclonal antibody BB27 recognizes CD101, a 140 kDa disulfide-bonded homodimeric protein expressed on activated T cells, and some other cell types, such as granulocytes and cells of the monocyte/macrophage lineage. HLDA V; WS Code T040
Regulatory Status:	RUO
Immunogen:	Human thymic clone B12
Species Reactivity:	Human
Application:	Flow Cytometry Immunoprecipitation Western Blotting Immunohistochemistry (frozen sections)
Purity:	> 95% (by SDS-PAGE)
Purification:	Purified by protein-A affinity chromatography
Concentration:	1 mg/ml
Storage Buffer:	Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.4
Storage / Stability:	Store at 2-8°C. Do not freeze. Do not use after expiration date stamped on vial label.
Expiration:	See vial label
Lot Number:	See vial label
Background:	CD101 is a type I transmembrane glycoprotein, which forms disulfide-linked homodimers. It is expressed on activated T cells, as well as on granulocytes, monocytes, dendritic cells or mucosal T cells. It plays a major role in the activation of T cells by skin dendritic cells. Function of CD101 has not been fully elucidated, but in mice its knock-out results in liver autoimmune disease induced by <i>Novosphingobium aromaticivorans</i> .
References:	*Leukocyte Typing V., Schlossman S. et al. (Eds.), Oxford University Press (1995). Bagot M, Martinel I, Charue D, Weill F, Boulland ML, Wechsler J, Freeman GJ, Bensussan A, Boumsell L: CD101 is expressed by skin dendritic cells. Role in T-lymphocyte activation. <i>Tissue Antigens</i> . 1997 Nov;50(5):439-48. *Bagot M, Martinel I, Charue D, Boulland ML, Wechsler J, Bensussan A, Boumsell L: Functional role of CD101 on skin dendritic cells. <i>Adv Exp Med Biol</i> . 1997;417:227-32. *Grassi F, Dezutter-Dambuyant C, McIlroy D, Jacquet C, Yoneda K, Imamura S, Boumsell L, Schmitt D, Autran B, Debré P, Hosmalin A: Monocyte-derived dendritic cells have a phenotype comparable to that of dermal dendritic cells and display ultrastructural granules distinct from Birbeck granules. <i>J Leukoc Biol</i> . 1998 Oct;64(4):484-93.

For laboratory research only, not for drug, diagnostic or other use.



Antibodies

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at www.exbio.cz.

For laboratory research only, not for drug, diagnostic or other use.

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic
Tel: +420 261 090 666 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz