



11-780-C025

## Monoclonal Antibody to CD111 Purified Antibody (0.025 mg)

<b>Clone:</b>	R1.302
<b>Isotype:</b>	Mouse IgG1
<b>Specificity:</b>	The mouse monoclonal antibody R1.302 recognizes CD111 (also known as Nectin 1), a 75 kDa type I transmembrane glycoprotein broadly expressed on endothelial cells, epithelial cells, neuronal cells, megakaryocytes, and CD34-positive stem cells.
<b>Regulatory Status:</b>	RUO
<b>Immunogen:</b>	NIH/3T3 cells transfected with human CD111
<b>Species Reactivity:</b>	Human
<b>Application:</b>	Flow Cytometry Immunoprecipitation Western Blotting Immunohistochemistry Immunocytochemistry
<b>Purity:</b>	> 95% (by SDS-PAGE)
<b>Purification:</b>	Purified by protein-A affinity chromatography
<b>Concentration:</b>	1 mg/ml
<b>Storage Buffer:</b>	Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.4
<b>Storage / Stability:</b>	Store at 2-8°C. Do not freeze. Do not use after expiration date stamped on vial label.
<b>Expiration:</b>	See vial label
<b>Lot Number:</b>	See vial label
<b>Background:</b>	CD111, also known as nectin-1, is a calcium-independent cell-cell adhesion transmembrane glycoprotein involved in organization of adherens junctions and tight junctions in epithelial and endothelial cells. It also serves as a target molecule for entry of herpes simplex virus (HSV-1, HSV-2) and pseudorabies virus (PRV) into epithelial and neuronal cells. CD111 is connected with actin cytoskeleton through afadin. Mutations in the gene for CD111 cause cleft lip and palate/ectodermal dysplasia 1 syndrome (CLPED1) as well as non-syndromic cleft lip with or without cleft palate (CL/P). Alternative splicing results in multiple transcript variants encoding proteins with distinct C-termini.

**For laboratory research only, not for drug, diagnostic or other use.**



**Antibodies**

**References:**

- \*Cocchi F, Lopez M, Dubreuil P, Campadelli Fiume G, Menotti L: Chimeric nectin1-poliiovirus receptor molecules identify a nectin1 region functional in herpes simplex virus entry. *J Virol.* 2001 Sep;75(17):7987-94.
- \*Cocchi F, Lopez M, Menotti L, Aoubala M, Dubreuil P, Campadelli-Fiume G: The V domain of herpesvirus Ig-like receptor (HIgR) contains a major functional region in herpes simplex virus-1 entry into cells and interacts physically with the viral glycoprotein D.
- \*Cocchi F, Lopez M, Dubreuil P, Campadelli Fiume G, Menotti L: Chimeric nectin1-poliiovirus receptor molecules identify a nectin1 region functional in herpes simplex virus entry. *J Virol.* 2001 Sep;75(17):7987-94.
- \*Reymond N, Garrido-Urbani S, Borg JP, Dubreuil P, Lopez M: PICK-1: a scaffold protein that interacts with Nectins and JAMs at cell junctions. *FEBS Lett.* 2005 Apr 11;579(10):2243-9.
- \*Yoon M, Spear PG: Disruption of adherens junctions liberates nectin-1 to serve as receptor for herpes simplex virus and pseudorabies virus entry. *J Virol.* 2002 Jul;76(14):7203-8.

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at [www.exbio.cz](http://www.exbio.cz).

**For laboratory research only, not for drug, diagnostic or other use.**

---

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic  
Tel: +420 261 090 666 | Fax: +420 261 090 660 | [orders@exbio.cz](mailto:orders@exbio.cz) | [www.exbio.cz](http://www.exbio.cz)