



11-788-C100

## Monoclonal Antibody to CD129 / IL-9R alpha Purified Antibody (0.1 mg)

<b>Clone:</b>	AH9R7
<b>Isotype:</b>	Mouse IgG2b
<b>Specificity:</b>	The mouse monoclonal antibody AH9R7 recognizes CD129 / IL-9R alpha, a 57 kDa type I transmembrane glycoprotein expressed at low levels by lymphocytes, blood cell progenitors, eosinophils, mast cells, epithelial cells, muscle cells and neurons.
<b>Regulatory Status:</b>	RUO
<b>Immunogen:</b>	Human CD129-transfected cell line
<b>Species Reactivity:</b>	Human
<b>Application:</b>	Flow Cytometry Application note: it is recommended to use bright fluorochromes or signal multiplying detection ELISA Functional Application blocking
<b>Purity:</b>	> 95% (by SDS-PAGE)
<b>Purification:</b>	Purified by protein-A affinity chromatography
<b>Concentration:</b>	1 mg/ml
<b>Storage Buffer:</b>	Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.4
<b>Storage / Stability:</b>	Store at 2-8°C. Do not freeze. Do not use after expiration date stamped on vial label.
<b>Expiration:</b>	See vial label
<b>Lot Number:</b>	See vial label
<b>Background:</b>	CD129 serves as the high affinity alpha subunit of IL-9 receptor. It associates with CD132, the common gamma chain shared by receptors of many different cytokines. CD129 is expressed at low levels by T and B cells, blood cell progenitors, eosinophils, mast cells, epithelial cells, muscle cells and neurons. Its signaling (through JAK/STAT pathways) results in proliferative and anti-apoptotic response, which is critical e.g. for intrathymic T cell development and survival of various cell types. The gene for CD129 is located at the pseudoautosomal regions of X and Y chromosomes and it may be related with the development of asthma.

**For laboratory research only, not for drug, diagnostic or other use.**



**Antibodies**

**References:**

De Smedt M, Verhasselt B, Kerre T, Vanhecke D, Naessens E, Leclercq G, Renauld JC, Van Snick J, Plum J: Signals from the IL-9 receptor are critical for the early stages of human intrathymic T cell development. *J Immunol.* 2000 Feb 15;164(4):1761-7.

\*Pilette C, Ouadrhiri Y, Van Snick J, Renauld JC, Staquet P, Vaerman JP, Sibille Y: IL-9 inhibits oxidative burst and TNF-alpha release in lipopolysaccharide-stimulated human monocytes through TGF-beta. *J Immunol.* 2002 Apr 15;168(8):4103-11.

Grasso L, Huang M, Sullivan CD, Messler CJ, Kiser MB, Dragwa CR, Holroyd KJ, Renauld JC, Levitt RC, Nicolaides NC: Molecular analysis of human interleukin-9 receptor transcripts in peripheral blood mononuclear cells. Identification of a splice variant encoding for a nonfunctional cell surface receptor. *J Biol Chem.* 1998 Sep 11;273(37):24016-24.

\*Pilette C, Ouadrhiri Y, Van Snick J, Renauld JC, Staquet P, Vaerman JP, Sibille Y: Oxidative burst in lipopolysaccharide-activated human alveolar macrophages is inhibited by interleukin-9. *Eur Respir J.* 2002 Nov;20(5):1198-205.

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at [www.exbio.cz](http://www.exbio.cz).

**For laboratory research only, not for drug, diagnostic or other use.**

---

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic  
Tel: +420 261 090 666 | Fax: +420 261 090 660 | [orders@exbio.cz](mailto:orders@exbio.cz) | [www.exbio.cz](http://www.exbio.cz)