



1A-158-T100

## Monoclonal Antibody to CD57 Allophycocyanin (APC) conjugated (100 tests)

<b>Clone:</b>	TB01
<b>Isotype:</b>	Mouse IgM
<b>Specificity:</b>	The mouse monoclonal antibody TB01 recognizes CD57, a carbohydrate antigen present mainly on NK cells, NK T cells, and in neural tissue. HLDA VI; WS Code NK16
<b>Regulatory Status:</b>	RUO
<b>Immunogen:</b>	A pool of neuroblastoma cell lines
<b>Species Reactivity:</b>	Human
<b>Preparation:</b>	The purified antibody is conjugated with cross-linked Allophycocyanin (APC) under optimum conditions. The conjugate is purified by size-exclusion chromatography and adjusted for direct use. No reconstitution is necessary.
<b>Storage Buffer:</b>	The reagent is provided in stabilizing phosphate buffered saline (PBS) solution containing 0.02% sodium azide.
<b>Storage / Stability:</b>	Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light. Do not use after expiration date stamped on vial label.
<b>Usage:</b>	The reagent is designed for Flow Cytometry analysis of human blood cells using 10 µl reagent / 100 µl of whole blood or 10 <sup>6</sup> cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.
<b>Expiration:</b>	See vial label
<b>Lot Number:</b>	See vial label
<b>Background:</b>	CD57, also known as HNK1 or Leu7, is a sulphated trisaccharide (3-O-sulfoglucuronic acid beta1-3 Gal beta1-4 GlcNAc) attached to several glycoproteins, including CD56, myelin glycoprotein PO, and neural cell adhesion molecule L1, as well as on glycolipids and chondroitin sulphate proteoglycans in the nervous system. It serves as a NK cell marker and it is expressed on well differentiated prostate cancers and uveal and cutaneous melanoma. CD57+ T cells are implicated as suppressors of T-cell responses.

**For laboratory research only, not for drug, diagnostic or other use.**

**Antibodies****References:**

- \*Lim SA, Kim TJ, Lee JE, Sonn CH, Kim K, Kim J, Choi JG, Choi IK, Yun CO, Kim JH, Yee C, Kumar V, Lee KM: Ex vivo expansion of highly cytotoxic human NK cells by cocultivation with irradiated tumor cells for adoptive immunotherapy. *Cancer Res.* 2013 Apr 15;73(8):2598-607.
- \*Cunha KS, Caruso AC, Faria PA, Silva LE, Fonseca EC, Geller M, de Moura-Neto RS, Lopes VS: Evaluation of Bcl-2, Bcl-x and cleaved caspase-3 in malignant peripheral nerve sheath tumors and neurofibromas. *An Acad Bras Cienc.* 2013;85(4):1497-511.
- \*Phadwal K, Alegre-Abarategui J, Watson AS, Pike L, Anbalagan S, Hammond EM, Wade-Martins R, McMichael A, Klenerman P, Simon AK: A novel method for autophagy detection in primary cells: impaired levels of macroautophagy in immunosenescent T cells. *Autophagy.* 2012 Apr;8(4):677-89.
- \*Marquardt N, Béziat V, Nyström S, Hengst J, Ivarsson MA, Kekäläinen E, Johansson H, Mjösberg J, Westgren M, Lankisch TO, Wedemeyer H, Ellis EC, Ljunggren HG, Michaëlsson J, Björkström NK: Cutting edge: identification and characterization of human intrahepatic CD49a+ NK cells. *J Immunol.* 2015 Mar 15;194(6):2467-71.
- \*Wangerin H, Kristiansen G, Schlomm T, Stephan C, Gunia S, Zimpfer A, Weichert W, Sauter G, Erbersdobler A: CD57 expression in incidental, clinically manifest, and metastatic carcinoma of the prostate. *Biomed Res Int.* 2014;2014:356427.
- \*Fernandez S, French MA, Price P: Immunosenescent CD57+CD4+ T-cells accumulate and contribute to interferon- $\gamma$  responses in HIV patients responding stably to ART. *Dis Markers.* 2011;31(6):337-42.

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at [www.exbio.cz](http://www.exbio.cz).

**For laboratory research only, not for drug, diagnostic or other use.**

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic  
Tel: +420 261 090 666 | Fax: +420 261 090 660 | [orders@exbio.cz](mailto:orders@exbio.cz) | [www.exbio.cz](http://www.exbio.cz)