



Antibodies

1A-314-T100

Monoclonal Antibody to CD177 Allophycocyanin (APC) conjugated (100 tests)

Clone:	MEM-166
Isotype:	Mouse IgG1
Specificity:	The antibody MEM-166 reacts with CD177 (Neutrophil specific antigen 1), a 60 kDa GPI-linked cell surface glycoprotein of uPAR family, expressed on granulocytes and in bone marrow early erythroblasts, megakaryocytes, promyelocytes and myelocytes. HLDA VI; WS Code M M17 HLDA VI; WS Code BP 309
Regulatory Status:	RUO
Immunogen:	Human granulocytes
Species Reactivity:	Human, Non-Human Primates
Preparation:	The purified antibody is conjugated with cross-linked Allophycocyanin (APC) under optimum conditions. The conjugate is purified by size-exclusion chromatography and adjusted for direct use. No reconstitution is necessary.
Storage Buffer:	The reagent is provided in stabilizing phosphate buffered saline (PBS) solution containing 15mM sodium azide.
Storage / Stability:	Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light. Do not use after expiration date stamped on vial label.
Usage:	The reagent is designed for Flow Cytometry analysis of human blood cells using 10 µl reagent / 100 µl of whole blood or 10 ⁶ cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.
Expiration:	See vial label
Lot Number:	See vial label
Background:	CD177 (NB1/HNA-2a and PRV-1 form) is a GPI-anchored glycoprotein present mainly on neutrophils. Its plasma membrane expression is increased during pregnancy and inflammation or after G-CSF application. Ligand of CD177 has been identified as CD31 (PECAM-1). CD177 participates in neutrophil transmigration and seems to be also a pro-proliferative molecule. The antibodies against CD177 can be involved in neonatal alloimmune neutropenia (NAN).

For laboratory research only, not for drug, diagnostic or other use.



Antibodies

References:

- *Leukocyte Typing VII., Mason D. et al. (Eds.), Oxford University Press (2002).
- *Stroncek DF, Caruccio L, Bettinotti M: CD177: A member of the Ly-6 gene superfamily involved with neutrophil proliferation and polycythemia vera. *J Transl Med.* 2004 Mar 29;2(1):8.
- *Mnjoyan Z, Li J, Afshar-Kharghan V: Expression of polycythemia rubra vera-1 decreases the dependency of cells on growth factors for proliferation. *Haematologica.* 2005 Mar;90(3):405-6.
- *Sachs UJ, Andrei-Selmer CL, Maniar A, Weiss T, Paddock C, Orlova VV, Choi EY, Newman PJ, Preissner KT, Chavakis T, Santoso S: The neutrophil-specific antigen CD177 is a counter-receptor for platelet endothelial cell adhesion molecule-1 (CD31). *J Biol Chem.* 2007 Aug 10;282(32):23603-12.
- *Bauer S, Abdgawad M, Gunnarsson L, Segelmark M, Tapper H, Hellmark T: Proteinase 3 and CD177 are expressed on the plasma membrane of the same subset of neutrophils. *J Leukoc Biol.* 2007 Feb;81(2):458-64.
- *Leukocyte Typing VI., Kishimoto T. et al. (Eds.), Garland Publishing Inc. (1997).

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at www.exbio.cz.

For laboratory research only, not for drug, diagnostic or other use.

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic
Tel: +420 261 090 666 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz