

1A-316-C100

Monoclonal Antibody to CD253 / TRAIL Allophycocyanin (APC) conjugated (0.1 mg)

Clone:	2E5
Isotype:	Mouse IgG1
Specificity:	The antibody 2E5 reacts with TRAIL (APO-2L), a 21 kDa cytotoxic protein, activator of rapid apoptosis in tumor cells. TRAIL is mainly expressed in spleen, lung, prostate and also in many other tissues.
Regulatory Status:	RUO
Immunogen:	Recombinant soluble fragment (aa 95-281) of human TRAIL.
Species Reactivity:	Human
Negative Species:	Mouse, Other not tested
Preparation:	The purified antibody is conjugated with cross-linked Allophycocyanin (APC) under optimum conditions. The conjugate is purified by size-exclusion chromatography.
Concentration:	0.1 mg/ml
Storage Buffer:	The reagent is provided in phosphate buffered saline (PBS) containing 15 mM sodium azide
Storage / Stability:	Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light. Do not use after expiration date stamped on vial label.
Usage:	The reagent is designed for Flow Cytometry analysis.
Expiration:	See vial label
Lot Number:	See vial label
Background:	Human CD253 / TRAIL (TNF-Related Apoptosis Inducing Ligand), also called Apo2, is a type II membrane protein from the TNF family. TRAIL is a cytotoxic protein which activates rapid apoptosis in tumor cells, but not in normal cells. TRAIL-induced apotosis, is achieved through binding to two dealth-signaling receptors, DR4 (CD261 / TRAIL-R1) and DR5 (CD262 / TRAIL-R2).
References:	*Plasilova M, Zivny J, Jelinek J, Neuwirtova R, Cermak J, Necas E, Andera L, Stopka T: TRAIL (Apo2L) suppresses growth of primary human leukemia and myelodysplasia progenitors. Leukemia. 2002 Jan;16(1):67-73. *Hyer ML, Croxton R, Krajewska M, Krajewski S, Kress CL, Lu M, Suh N, Sporn MB, Cryns VL, Zapata JM, Reed JC: Synthetic triterpenoids cooperate with tumor necrosis factor-related apoptosis-inducing ligand to induce apoptosis of breast cancer cells. Cancer Res. 2005 Jun 1;65(11):4799-808.

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at www.exbio.cz.

For laboratory research only, not for drug, diagnostic or other use.

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic Tel: +420 261 090 666 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz