



1A-506-T025

## Monoclonal Antibody to CD22 Allophycocyanin (APC) conjugated (25 tests)

<b>Clone:</b>	IS7
<b>Isotype:</b>	Mouse IgG1
<b>Specificity:</b>	<p>The antibody IS7 reacts with CD22 (BL-CAM), a 130 kDa type I transmembrane glycoprotein (immunoglobulin superfamily) expressed in the cytoplasm of pro-B and pre-B lymphocytes, and on the surface of mature and activated B lymphocytes; it is lost on plasma cells, peripheral blood T lymphocytes, granulocytes and monocytes.</p> <p>HLDA IV; WS Code B 227 HLDA V; WS Code B CD22.8</p>
<b>Regulatory Status:</b>	RUO
<b>Immunogen:</b>	human cell line Reh
<b>Species Reactivity:</b>	Human
<b>Preparation:</b>	The purified antibody is conjugated with cross-linked Allophycocyanin (APC) under optimum conditions. The conjugate is purified by size-exclusion chromatography and adjusted for direct use. No reconstitution is necessary.
<b>Storage Buffer:</b>	The reagent is provided in stabilizing phosphate buffered saline (PBS) solution containing 15mM sodium azide.
<b>Storage / Stability:</b>	Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light. Do not use after expiration date stamped on vial label.
<b>Usage:</b>	<p>The reagent is designed for Flow Cytometry analysis of human blood cells using 10 µl reagent / 100 µl of whole blood or 10<sup>6</sup> cells in a suspension.</p> <p>The content of a vial (0.25 ml) is sufficient for 25 tests.</p>
<b>Expiration:</b>	See vial label
<b>Lot Number:</b>	See vial label
<b>Background:</b>	<p>CD22, also known as Siglec-2 (sialic acid-binding immunoglobulin-like lectin-2) is a transmembrane glycoprotein binding alpha2,6-linked sialic acid-bearing ligands. Intracellular domain of CD22 recruits protein tyrosine phosphatase SHP-1 through the immunoreceptor tyrosine-based inhibitory motifs (ITIMs), thus setting a threshold for B cell receptor-mediated activation. CD22 also regulates B-cell response by involvement in controlling the CD19/CD21-Src-family protein tyrosine kinase amplification pathway and CD40 signaling. CD22 exhibits hallmarks of clathrin-mediated endocytic pathway.</p>
<b>References:</b>	<p>*Tedder TF, Poe JC, Haas KM: CD22: A Multifunctional Receptor That Regulates B Lymphocyte Survival and Signal Transduction. <i>Adv Immunol.</i> 2005;88:1-50.</p> <p>*Tateno H, Li H, Schur MJ, Bovin N, Crocker PR, Wakarchuk WW, Paulson JC: Distinct endocytic mechanisms of CD22 (Siglec-2) and Siglec-F reflect roles in cell signaling and innate immunity. <i>Mol Cell Biol.</i> 2007 Aug;27(16):5699-710.</p> <p>*Walker JA, Smith KG: CD22: an inhibitory enigma. <i>Immunology.</i> 2007 Dec 7</p> <p>*Leukocyte Typing IV., Knapp W. et al. (Eds.), Oxford University Press (1989).</p> <p>*Leukocyte Typing V., Schlossman S. et al. (Eds.), Oxford University Press (1995).</p>

**For laboratory research only, not for drug, diagnostic or other use.**



**Antibodies**

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at [www.exbio.cz](http://www.exbio.cz).

**For laboratory research only, not for drug, diagnostic or other use.**

---

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic  
Tel: +420 261 090 666 | Fax: +420 261 090 660 | [orders@exbio.cz](mailto:orders@exbio.cz) | [www.exbio.cz](http://www.exbio.cz)