



Antibodies

1A-559-T100

Monoclonal Antibody to CD82 Allophycocyanin (APC) conjugated (100 tests)

Clone:	C33
Isotype:	Mouse IgG2a
Specificity:	The antibody C33 reacts with CD82, a widely expressed cell surface protein of the tetraspanin family. CD82 is also found in endosome/lysosome compartments.
Regulatory Status:	RUO
Immunogen:	C91/PL (human HTLV-1+ T cell line)
Species Reactivity:	Human, Other not tested
Preparation:	The purified antibody is conjugated with cross-linked Allophycocyanin (APC) under optimum conditions. The conjugate is purified by size-exclusion chromatography and adjusted for direct use. No reconstitution is necessary.
Storage Buffer:	The reagent is provided in stabilizing phosphate buffered saline (PBS) solution containing 15mM sodium azide.
Storage / Stability:	Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light. Do not use after expiration date stamped on vial label.
Usage:	The reagent is designed for Flow Cytometry analysis of human blood cells using 10 µl reagent / 100 µl of whole blood or 10 ⁶ cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.
Expiration:	See vial label
Lot Number:	See vial label
Background:	CD82 (KAI1), a member of the tetraspanin family, forms complexes with other tetraspanin proteins, integrins, coreceptors, MHC class I and II molecules. These complexes influence adhesion, morphology, activation, proliferation and differentiation of B, T and other cells. CD82 regulates cytoskeleton rearrangement and may participate in the turnover of the tetraspanin complex members. Besides in the plasma membrane, CD82 is localized also in endosome/lysosome compartments. Tumour-suppressive roles of CD82 have been demonstrated.

For laboratory research only, not for drug, diagnostic or other use.

**Antibodies****References:**

- *Fukudome K, Furuse M, Imai T, Nishimura M, Takagi S, Hinuma Y, Yoshie O: Identification of membrane antigen C33 recognized by monoclonal antibodies inhibitory to human T-cell leukemia virus type 1 (HTLV-1)-induced syncytium formation: altered glycosylation of C33 antigen in HTLV-1-positive T cells. *J Virol.* 1992 Mar;66(3):1394-401.
- *Imai T, Yoshie O: C33 antigen and M38 antigen recognized by monoclonal antibodies inhibitory to syncytium formation by human T cell leukemia virus type 1 are both members of the transmembrane 4 superfamily and associate with each other and with CD4 or CD8 in T cells. *J Immunol.* 1993 Dec 1;151(11):6470-81.
- *Imai T, Kakizaki M, Nishimura M, Yoshie O: Molecular analyses of the association of CD4 with two members of the transmembrane 4 superfamily, CD81 and CD82. *J Immunol.* 1995 Aug 1;155(3):1229-39.
- *Escola JM, Kleijmeer MJ, Stoorvogel W, Griffith JM, Yoshie O, Geuze HJ: Selective enrichment of tetraspan proteins on the internal vesicles of multivesicular endosomes and on exosomes secreted by human B-lymphocytes. *J Biol Chem.* 1998 Aug 7;273(32):20121-7.
- *Ueda T, Ichikawa T, Tamaru J, Mikata A, Akakura K, Akimoto S, Imai T, Yoshie O, Shiraishi T, Yatani R, Ito H, Shimazaki J: Expression of the KAI1 protein in benign prostatic hyperplasia and prostate cancer. *Am J Pathol.* 1996 Nov;149(5):1435-40.
- *Schatzmaier P, Supper V, Göschl L, Zwirzitz A, Eckerstorfer P, Ellmeier W, Huppa JB, Stockinger H: Rapid multiplex analysis of lipid raft components with single-cell resolution. *Sci Signal.* 2015 Sep 22;8(395):rs11

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at www.exbio.cz.

For laboratory research only, not for drug, diagnostic or other use.

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic
Tel: +420 261 090 666 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz