



1A-581-T025

Monoclonal Antibody to CD326 / EpCAM Allophycocyanin (APC) conjugated (25 tests)

Clone:	VU-1D9
Isotype:	Mouse IgG1
Specificity:	The mouse monoclonal antibody VU-1D9 recognizes an epitope within EGF-like domain I of CD326 / EpCAM, a marker of epithelial lineages. This antibody strongly stains various normal epithelial cells and carcinomas.
Regulatory Status:	RUO
Immunogen:	Small cell lung carcinoma cell line H69.
Species Reactivity:	Human, Other species Not tested
Preparation:	The purified antibody is conjugated with cross-linked Allophycocyanin (APC) under optimum conditions. The conjugate is purified by size-exclusion chromatography and adjusted for direct use. No reconstitution is necessary.
Storage Buffer:	The reagent is provided in stabilizing phosphate buffered saline (PBS) solution containing 15mM sodium azide.
Storage / Stability:	Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light. Do not use after expiration date stamped on vial label.
Usage:	The reagent is designed for Flow Cytometry analysis of human blood cells using 10 µl reagent / 100 µl of whole blood or 10 ⁶ cells in a suspension. The content of a vial (0.25 ml) is sufficient for 25 tests.
Expiration:	See vial label
Lot Number:	See vial label
Background:	CD326 / EpCAM (also known as ESA, EGP40, EGP-2, KSA1/4, CO17-1A, GA733-2, MOC31, Ber-EP4) is a 40 kDa transmembrane glycoprotein serving as adhesion molecule in the basolateral membranes in a variety of epithelial cells. CD326 mediates calcium-independent homotypic cell-cell adhesions. CD326 over-expression has been detected in many epithelial tumours and is often associated with bad prognosis. It has been used for diagnostics of (pre-) malignancies at early stages.
References:	*Tsubura A, Senzaki H, Sasaki M, Hilgers J, Morii S: Immunohistochemical demonstration of breast-derived and/or carcinoma-associated glycoproteins in normal skin appendages and their tumors. J Cutan Pathol. 1992 Feb;19(1):73-9. *Ogura E, Senzaki H, Yoshizawa K, Hioki K, Tsubura A: Immunohistochemical localization of epithelial glycoprotein EGP-2 and carcinoembryonic antigen in normal colonic mucosa and colorectal tumors. Anticancer Res. 1998 Sep-Oct;18(5B):3669-75. *Li G, Passebosch-Faure K, Lambert C, Gentil-Perret A, Blanc F, Oosterwijk E, Mosnier JF, Genin C, Tostain J: Flow cytometric analysis of antigen expression in malignant and normal renal cells. Anticancer Res. 2000 Jul-Aug;20(4):2773-8. *Winter MJ, Nagtegaal ID, van Krieken JH, Litvinov SV: The epithelial cell adhesion molecule (Ep-CAM) as a morphoregulatory molecule is a tool in surgical pathology. Am J Pathol. 2003 Dec;163(6):2139-48. *Brunner A, Prelog M, Verdorfer I, Tzankov A, Mikuz G, Ensinger C: EpCAM is predominantly expressed in high grade and advanced stage urothelial carcinoma of the bladder. J Clin Pathol. 2008 Mar;61(3):307-10.

For laboratory research only, not for drug, diagnostic or other use.



Antibodies

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at www.exbio.cz.

For laboratory research only, not for drug, diagnostic or other use.

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic
Tel: +420 261 090 666 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz