



Antibodies

1A-588-T025

## Monoclonal Antibody to CD324 / E-Cadherin Allophycocyanin (APC) conjugated (25 tests)

|                             |   |
|-----------------------------|---|
| <b>Clone:</b>               | 67A4  |
| <b>Isotype:</b>             | Mouse IgG1  |
| <b>Specificity:</b>         | The mouse monoclonal antibody 67A4 recognizes CD324 / E-cadherin, an approximately 100 kDa epithelial cell adhesion molecule, whose detection is important for determination of invasive potential of epithelial neoplasms.<br>HLDA VIII  |
| <b>Regulatory Status:</b>   | RUO   |
| <b>Immunogen:</b>           | T-47D cells   |
| <b>Species Reactivity:</b>  | Human   |
| <b>Preparation:</b>         | The purified antibody is conjugated with cross-linked Allophycocyanin (APC) under optimum conditions. The conjugate is purified by size-exclusion chromatography and adjusted for direct use. No reconstitution is necessary.   |
| <b>Storage Buffer:</b>      | The reagent is provided in stabilizing phosphate buffered saline (PBS) solution containing 15mM sodium azide.   |
| <b>Storage / Stability:</b> | Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light. Do not use after expiration date stamped on vial label.   |
| <b>Usage:</b>               | The reagent is designed for Flow Cytometry analysis of human blood cells using 10 µl reagent / 100 µl of whole blood or 10 <sup>6</sup> cells in a suspension.<br>The content of a vial (0.25 ml) is sufficient for 25 tests.   |
| <b>Expiration:</b>          | See vial label  |
| <b>Lot Number:</b>          | See vial label  |
| <b>Background:</b>          | CD324 / E-cadherin is an epithelial cell surface molecule, which provides calcium-dependent homophilic interactions with E-cadherin of another cell. These interactions take part in morphogenetic programs controlling the maintenance of the structural and functional integrity of epithelia and affect invasive potential of epithelial neoplasms. CD324 / E-cadherin is implicated in cell growth and differentiation, cell recognition, and sorting during developmental morphogenesis, as well as in aggregation-dependent cell survival. CD324 / E-cadherin-mediated cell adhesion system is highly regulated from inside the cell by a number of intracellular signaling pathways. |

**For laboratory research only, not for drug, diagnostic or other use.**

**Antibodies****References:**

- \*Takeichi M: Cadherin cell adhesion receptors as a morphogenetic regulator. *Science*. 1991 Mar 22;251(5000):1451-5.
- \*Pece S, Chiariello M, Murga C, Gutkind JS: Activation of the protein kinase Akt/PKB by the formation of E-cadherin-mediated cell-cell junctions. Evidence for the association of phosphatidylinositol 3-kinase with the E-cadherin adhesion complex. *J Biol Chem*. 1999 Jul 2;274(27):19347-51.
- \*Pece S, Gutkind JS: Signaling from E-cadherins to the MAPK pathway by the recruitment and activation of epidermal growth factor receptors upon cell-cell contact formation. *J Biol Chem*. 2000 Dec 29;275(52):41227-33.
- \*Armeanu S, Bühring HJ, Reuss-Borst M, Müller CA, Klein G: E-cadherin is functionally involved in the maturation of the erythroid lineage. *J Cell Biol*. 1995 Oct;131(1):243-9.
- \*Bühring HJ, Müller T, Herbst R, Cole S, Rappold I, Schuller W, Zhu X, Fritzscher U, Faul C, Armeanu S, Ullrich A, Klein G, Schmidt H: The adhesion molecule E-cadherin and a surface antigen recognized by the antibody 9C4 are selectively expressed on erythroid cells of defined maturational stages. *Leukemia*. 1996 Jan;10(1):106-16.
- \*Novak N, Kraft S, Haberstok J, Geiger E, Allam P, Bieber T: A reducing microenvironment leads to the generation of FcepsilonRIhigh inflammatory dendritic epidermal cells (IDEC). *J Invest Dermatol*. 2002 Oct;119(4):842-9.
- \*Servet-Delprat C, Vidalain PO, Bausinger H, Manié S, Le Deist F, Azocar O, Hanau D, Fischer A, Rabourdin-Combe C: Measles virus induces abnormal differentiation of CD40 ligand-activated human dendritic cells. *J Immunol*. 2000 Feb 15;164(4):1753-60.
- \*Furio L, Guezennec A, Ducarre B, Guesnet J, Peguet-Navarro J: Differential effects of allergens and irritants on early differentiating monocyte-derived dendritic cells. *Eur J Dermatol*. 2008 Mar-Apr;18(2):141-7.
- \*Robertson H, Ali S, McDonnell BJ, Burt AD, Kirby JA: Chronic renal allograft dysfunction: the role of T cell-mediated tubular epithelial to mesenchymal cell transition. *J Am Soc Nephrol*. 2004 Feb;15(2):390-7.
- \*Kutlesa S, Wessels JT, Speiser A, Steiert I, Müller CA, Klein G: E-cadherin-mediated interactions of thymic epithelial cells with CD103+ thymocytes lead to enhanced thymocyte cell proliferation. *J Cell Sci*. 2002 Dec 1;115(Pt 23):4505-15.
- \*Caberg JH, Hubert PM, Begon DY, Herfs MF, Roncarati PJ, Boniver JJ, Delvenne PO: Silencing of E7 oncogene restores functional E-cadherin expression in human papillomavirus 16-transformed keratinocytes. *Carcinogenesis*. 2008 Jul;29(7):1441-7.
- \*Lin JC, Liao SK, Lee EH, Hung MS, Sayion Y, Chen HC, Kang CC, Huang LS, Cherng JM. Molecular events associated with epithelial to mesenchymal transition of nasopharyngeal carcinoma cells in the absence of Epstein-Barr virus genome. *J Biomed Sci*. 2009 Nov 24;16:105.
- \*And many other.

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at [www.exbio.cz](http://www.exbio.cz).

**For laboratory research only, not for drug, diagnostic or other use.**

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic  
Tel: +420 261 090 666 | Fax: +420 261 090 660 | [orders@exbio.cz](mailto:orders@exbio.cz) | [www.exbio.cz](http://www.exbio.cz)