

1A-601-C100

Monoclonal Antibody to FoxP3 Allophycocyanin (APC) conjugated (0.1 mg)

Clone:	3G3
Isotype:	Mouse IgG1
Specificity:	The mouse monoclonal antibody 3G3 recognizes N-terminal region of FoxP3, a 47-55 kDa transcription factor, which is the master regulator in the development and function of regulatory T cells.
Regulatory Status:	RUO
Immunogen:	Full-length His-tagged recombinant murine FoxP3
Species Reactivity:	Human, Mouse
Preparation:	The purified antibody is conjugated with cross-linked Allophycocyanin (APC) under optimum conditions. The conjugate is purified by size-exclusion chromatography.
Concentration:	0.5 mg/ml
Storage Buffer:	Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.4
Storage / Stability:	Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light. Do not use after expiration date stamped on vial label.
Usage:	The reagent is designed for Flow Cytometry analysis.
Expiration:	See vial label
Lot Number:	See vial label
Background:	FoxP3 (Forkhead box protein 3), a highly conserved forkhead/winged-helix transcription factor, plays a crucial role in maintaining immune homeostasis by governing the development and function of regulatory T cells. It is constitutively expressed at high level in CD25 ⁺ CD4 ⁺ Treg cells and at low level in a CD25 ⁻ CD4 ⁺ Treg cell subset. Defects in gene encoding FoxP3 protein cause the scurfy phenotype in mice, and in human the IPEX syndrome (immune dysfunction, polyendocrinopathy, enteropathy, X-linked syndrome), also known as X-linked autoimmunity-allergic dysregulation (XLAAD) syndrome.

For laboratory research only, not for drug, diagnostic or other use.



Antibodies

- References:**
- *Bettini M, Vignali DA: Regulatory T cells and inhibitory cytokines in autoimmunity. *Curr Opin Immunol.* 2009 Dec;21(6):612-8.
 - *Barnes MJ, Powrie F: Regulatory T cells reinforce intestinal homeostasis. *Immunity.* 2009 Sep 18;31(3):401-11.
 - *Kuhn A, Beissert S, Krammer PH: CD4(+)CD25 (+) regulatory T cells in human lupus erythematosus. *Arch Dermatol Res.* 2009 Jan;301(1):71-81.
 - *Elkord E: Novel therapeutic strategies by regulatory T cells in allergy. *Chem Immunol Allergy.* 2008;94:150-7.
 - *Lal G, Bromberg JS: Epigenetic mechanisms of regulation of Foxp3 expression. *Blood.* 2009 Oct 29;114(18):3727-35.
 - *Gavin MA, Torgerson TR, Houston E, DeRoos P, Ho WY, Stray-Pedersen A, Ocheltree EL, Greenberg PD, Ochs HD, Rudensky AY: Single-cell analysis of normal and FOXP3-mutant human T cells: FOXP3 expression without regulatory T cell development. *Proc Natl Acad Sci U S A.* 2006 Apr 25;103(17):6659-64.
 - *Law JP, Hirschhorn DF, Owen RE, Biswas HH, Norris PJ, Lanteri MC: The importance of Foxp3 antibody and fixation/permeabilization buffer combinations in identifying CD4+CD25+Foxp3+ regulatory T cells. *Cytometry A.* 2009 Dec;75(12):1040-50.

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at www.exbio.cz.

For laboratory research only, not for drug, diagnostic or other use.

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic
Tel: +420 261 090 666 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz