

1A-715-C100

Monoclonal Antibody to CD45R (human, mouse) Allophycocyanin (APC) conjugated (0.1 mg)

Clone:	RA3-6B2
Isotype:	Rat IgG2a
Specificity:	The rat monoclonal antibody RA3-6B2 recognizes an epitope on CD45R, which is expressed at all developmental stages of B cells, including activated B cells, but also on subsets of NK and T cells. T cells detected by this antibody are supposed to be in activated state.
Regulatory Status:	RUO
Immunogen:	Abelson murine leukemia virus-induced pre-B tumor cells
Species Reactivity:	Human, Mouse, Feline (Cat)
Preparation:	The purified antibody is conjugated with cross-linked Allophycocyanin (APC) under optimum conditions. The conjugate is purified by size-exclusion chromatography.
Concentration:	0.5 mg/ml
Storage Buffer:	Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.4
Storage / Stability:	Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light. Do not use after expiration date stamped on vial label.
Usage:	The reagent is designed for Flow Cytometry analysis.
Expiration:	See vial label
Lot Number:	See vial label
Background:	CD45R, also known as B220, is a receptor-type protein tyrosine phosphatase glycoprotein. It is crucial in lymphocyte development and antigen signaling, serving as an important regulator of Src-family kinases, promotes cell survival by modulating integrin-mediated signal transduction pathway and is also involved in DNA fragmentation during apoptosis. CD45R expression also identifies a subset of murine bone marrow cells able to form osteoclast-like cells.

For laboratory research only, not for drug, diagnostic or other use.

**Antibodies**

- References:**
- *Watanabe Y, Akaike T: Activation signal induces the expression of B cell-specific CD45R epitope (6B2) on murine T cells. *Scand J Immunol.* 1994 May;39(5):419-25.
 - *Tsurui H, Nishimura H, Hattori S, Hirose S, Okumura K, Shirai T: Seven-color fluorescence imaging of tissue samples based on Fourier spectroscopy and singular value decomposition. *J Histochem Cytochem.* 2000 May;48(5):653-62.
 - *Rada C, Ehrenstein MR, Neuberger MS, Milstein C: Hot spot focusing of somatic hypermutation in MSH2-deficient mice suggests two stages of mutational targeting. *Immunity.* 1998 Jul;9(1):135-41.
 - *Rothäusler K, Baumgarth N: Evaluation of intranuclear BrdU detection procedures for use in multicolor flow cytometry. *Cytometry A.* 2006 Apr;69(4):249-59.
 - *Montecino-Rodriguez E, Leathers H, Dorshkind K: Identification of a B-1 B cell-specified progenitor. *Nat Immunol.* 2006 Mar;7(3):293-301.
 - *Cappione AJ, Pugh-Bernard AE, Anolik JH, Sanz I: Lupus IgG VH4.34 antibodies bind to a 220-kDa glycoform of CD45/B220 on the surface of human B lymphocytes. *J Immunol.* 2004 Apr 1;172(7):4298-307.
 - *Lochhead RB, Sonderegger FL, Ma Y, Brewster JE, Cornwall D, Maylor-Hagen H, Miller JC, Zachary JF, Weis JH, Weis JJ: Endothelial cells and fibroblasts amplify the arthritogenic type I IFN response in murine Lyme disease and are major sources of chemokines in *Borrelia burgdorferi*-infected joint tissue. *J Immunol.* 2012 Sep 1;189(5):2488-501

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at www.exbio.cz.

For laboratory research only, not for drug, diagnostic or other use.

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic
Tel: +420 261 090 666 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz