



1B-233-C100

Monoclonal Antibody to CD59 Biotin conjugated (0.1 mg)

Clone:	MEM-43
Isotype:	Mouse IgG2a
Specificity:	<p>The antibody MEM-43 reacts with well defined epitope (W40, R-53) on CD59 (Protectin), an 18-20 kDa glycosylphosphatidylinositol (GPI)-anchored glycoprotein expressed on all hematopoietic cells; it is widely present on cells in all tissues. This antibody does not compete with MEM-43/5.</p> <p>HLDA IV; WS Code NL 705 HLDA V; WS Code AS S013 HLDA V; WS Code BP BP345 HLDA V; WS Code T T-103</p>
Regulatory Status:	RUO
Immunogen:	Thymocytes and T lymphocytes
Species Reactivity:	Human
Preparation:	The purified antibody is conjugated with Biotin-LC-NHS under optimum conditions. The reagent is free of unconjugated biotin.
Concentration:	1 mg/ml
Storage Buffer:	Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.4
Storage / Stability:	Store at 2-8°C. Do not freeze. Do not use after expiration date stamped on vial label.
Usage:	<p>Biotinylated antibody is designed for indirect immunofluorescence analysis by Flow Cytometry.</p> <p>Suggested working dilution is 5 microgram/ml. Indicated dilution is recommended starting point for use of this product. Working concentrations should be determined by the investigator.</p>
Expiration:	See vial label
Lot Number:	See vial label
Background:	<p>CD59 (Protectin) is a small (18-20 kDa) GPI-anchored ubiquitously expressed inhibitor of the membrane attack complex (MAC). It is thus the key regulator that preserves the autologous cells from terminal effector mechanism of the complement cascade. CD59 associates with C5b-8 complex and thereby counteracts appropriate formation of cytolytic pore within the plasma membrane. CD59 is also an low-affinity ligand of human CD2 and causes T cell costimulation.</p>

For laboratory research only, not for drug, diagnostic or other use.

**Antibodies****References:**

- *Meri S, Morgan BP, Davies A, Daniels RH, Olavesen MG, Waldmann H, Lachmann PJ: Human protectin (CD59), an 18,000-20,000 MW complement lysis restricting factor, inhibits C5b-8 catalysed insertion of C9 into lipid bilayers. *Immunology*. 1990 Sep;71(1):1-9.
- *Rooney IA, Davies A, Griffiths D, Williams JD, Davies M, Meri S, Lachmann PJ, Morgan BP: The complement-inhibiting protein, protectin (CD59 antigen), is present and functionally active on glomerular epithelial cells. *Clin Exp Immunol*. 1991 Feb;83(2):251-6.
- *Menu E, Tsai BC, Bothwell AL, Sims PJ, Bierer BE: CD59 costimulation of T cell activation. CD58 dependence and requirement for glycosylation. *J Immunol*. 1994 Sep 15;153(6):2444-56.
- *Baalasubramanian S, Harris CL, Donev RM, Mizuno M, Omidvar N, Song WC, Morgan BP: CD59a is the primary regulator of membrane attack complex assembly in the mouse. *J Immunol*. 2004 Sep 15;173(6):3684-92.
- *Horejsi V, Hilgert I, Kristofova H, Bazil V, Bukovsky A, Kulhankova J: Monoclonal antibodies against human leucocyte antigens. I. Antibodies against beta-2-microglobulin, immunoglobulin kappa light chains, HLA-DR-like antigens, T8 antigen, T1 antigen, a monocyte antigen, and a pan-leucocyte antigen. *Folia Biol (Praha)*. 1986;32(1):12-25. (original description of MEM-43 antigen)
- *IMPORTANT ARTICLE: Robert Sutherland D, Keeney M, Illingworth A: Practical guidelines for the high-sensitivity detection and monitoring of paroxysmal nocturnal hemoglobinuria (PNH) clones by flow cytometry. *Cytometry B Clin Cytom*. 2012 Apr 12. doi: 10.1002/cyto.b.21023. [Epub ahead of print] Note: This article recommends PE-conjugated MEM-43 as a good reagent for red blood cell analysis of PNH (Paroxysmal Nocturnal Hemoglobinuria) by flow cytometry.
- *Leukocyte Typing IV., Knapp W. et al. (Eds.), Oxford University Press (1989).
- *Leukocyte Typing V., Schlossman S. et al. (Eds.), Oxford University Press (1995).
- *Forsberg UH, Bazil V, Stefanova I, Schroder J: Gene for human CD59 (likely Ly-6 homologue) is located on the short arm of chromosome 11. *Immunogenetics*. 1989;30(3):188-93.
- *Stefanova I, Hilgert I, Kristofova H, Brown R, Low MG, Horejsi V: Characterization of a broadly expressed human leucocyte surface antigen MEM-43 anchored in membrane through phosphatidylinositol. *Mol Immunol*. 1989 Feb;26(2):153-61.
- *Stefanova I, Horejsi V, Ansotegui IJ, Knapp W, Stockinger H: GPI-anchored cell-surface molecules complexed to protein tyrosine kinases. *Science*. 1991 Nov 15;254(5034):1016-9.
- *Cinek T, Horejsi V: The nature of large noncovalent complexes containing glycosyl-phosphatidylinositol-anchored membrane glycoproteins and protein tyrosine kinases. *J Immunol*. 1992 Oct 1;149(7):2262-70.
- *Bodian DL, Davis SJ, Morgan BP, Rushmere NK: Mutational analysis of the active site and antibody epitopes of the complement-inhibitory glycoprotein, CD59. *J Exp Med*. 1997 Feb 3;185(3):507-16.
- Cebecauer M, Cerny J, Horejsi V: Incorporation of leucocyte GPI-anchored proteins and protein tyrosine kinases into lipid-rich membrane domains of COS-7 cells. *Biochem Biophys Res Commun*. 1998 Feb 24;243(3):706-10.
- *Ilangumaran S, Briol A, Hoessli DC: CD44 selectively associates with active Src family protein tyrosine kinases Lck and Fyn in glycosphingolipid-rich plasma membrane domains of human peripheral blood lymphocytes. *Blood*. 1998 May 15;91(10):3901-8.
- *Omidvar N, Wang EC, Brennan P, Longhi MP, Smith RA, Morgan BP: Expression of glycosylphosphatidylinositol-anchored CD59 on target cells enhances human NK cell-mediated cytotoxicity. *J Immunol*. 2006 Mar 1;176(5):2915-23.
- *And many other.

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at www.exbio.cz.

For laboratory research only, not for drug, diagnostic or other use.

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic
Tel: +420 261 090 666 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz