

1B-265-C025

Monoclonal Antibody to CD22 Biotin conjugated (0.025 mg)

Clone:	MEM-01
Isotype:	Mouse IgG1
Specificity:	<p>The antibody MEM-01 reacts with CD22 (BL-CAM), a 130 kDa type I transmembrane glycoprotein (immunoglobulin superfamily) expressed in the cytoplasm of pro-B and pre-B lymphocytes, and on the surface of mature and activated B lymphocytes; it is lost on plasma cells, peripheral blood T lymphocytes, granulocytes and monocytes.</p> <p>The antibody MEM-01 cross-blocks the antibody OTH228 that recognizes uniquely epitope "E"; it does not cross-block antibodies RFB-4, CLB22/1 and CLB-BLy1.</p>
Regulatory Status:	RUO
Immunogen:	Raji Burkitt's lymphoma cell line
Species Reactivity:	Human, Non-Human Primates
Preparation:	The purified antibody is conjugated with Biotin-LC-NHS under optimum conditions. The reagent is free of unconjugated biotin.
Concentration:	1 mg/ml
Storage Buffer:	Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.4
Storage / Stability:	Store at 2-8°C. Do not freeze. Do not use after expiration date stamped on vial label.
Usage:	<p>Biotinylated antibody is designed for indirect immunofluorescence analysis by Flow Cytometry.</p> <p>Suggested working dilution is 1:500. Indicated dilution is recommended starting point for use of this product. Working concentrations should be determined by the investigator.</p>
Expiration:	See vial label
Lot Number:	See vial label
Background:	<p>CD22, also known as Siglec-2 (sialic acid-binding immunoglobulin-like lectin-2) is a transmembrane glycoprotein binding alpha2,6-linked sialic acid-bearing ligands. Intracellular domain of CD22 recruits protein tyrosine phosphatase SHP-1 through the immunoreceptor tyrosine-based inhibitory motifs (ITIMs), thus setting a threshold for B cell receptor-mediated activation. CD22 also regulates B-cell response by involvement in controlling the CD19/CD21-Src-family protein tyrosine kinase amplification pathway and CD40 signaling. CD22 exhibits hallmarks of clathrin-mediated endocytic pathway.</p>
References:	<p>*Tedder TF, Poe JC, Haas KM: CD22: A Multifunctional Receptor That Regulates B Lymphocyte Survival and Signal Transduction. <i>Adv Immunol.</i> 2005;88:1-50.</p> <p>*Tateno H, Li H, Schur MJ, Bovin N, Crocker PR, Wakarchuk WW, Paulson JC: Distinct endocytic mechanisms of CD22 (Siglec-2) and Siglec-F reflect roles in cell signaling and innate immunity. <i>Mol Cell Biol.</i> 2007 Aug;27(16):5699-710.</p> <p>*Walker JA, Smith KG: CD22: an inhibitory enigma. <i>Immunology.</i> 2007 Dec 7</p>

For laboratory research only, not for drug, diagnostic or other use.



Antibodies

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at www.exbio.cz.

For laboratory research only, not for drug, diagnostic or other use.

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic
Tel: +420 261 090 666 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz