



1B-558-C100

## Monoclonal Antibody to CD81 Biotin conjugated (0.1 mg)

<b>Clone:</b>	M38
<b>Isotype:</b>	Mouse IgG1
<b>Specificity:</b>	The antibody M38 reacts with CD81, a 25 kDa member of the tetraspanin family, expressed on majority of cells.
<b>Regulatory Status:</b>	RUO
<b>Immunogen:</b>	MOLT-4 (human T-ALL cell line)
<b>Species Reactivity:</b>	Human, Feline (Cat), Rabbit
<b>Preparation:</b>	The purified antibody is conjugated with Biotin-LC-NHS under optimum conditions. The reagent is free of unconjugated biotin.
<b>Concentration:</b>	1 mg/ml
<b>Storage Buffer:</b>	Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.4
<b>Storage / Stability:</b>	Store at 2-8°C. Do not freeze. Do not use after expiration date stamped on vial label.
<b>Usage:</b>	Biotinylated antibody is designed for indirect immunofluorescence analysis by Flow Cytometry.
<b>Expiration:</b>	See vial label
<b>Lot Number:</b>	See vial label
<b>Background:</b>	CD81 (TAPA-1), a member of the tetraspanin family, is expressed on virtually all nucleated cells, but above all on germinal center B cells. CD81 forms complexes with other tetraspanin proteins, integrins, coreceptors, MHC class I and II molecules, and influences adhesion, morphology, activation, proliferation and differentiation of B, T and other cells; e.g. in muscles CD81 promotes cell fusion and myotube maintenance. CD81 has been also identified as a receptor for the hepatitis C virus.

**For laboratory research only, not for drug, diagnostic or other use.**

**Antibodies****References:**

\*Fukudome K, Furuse M, Imai T, Nishimura M, Takagi S, Hinuma Y, Yoshie O: Identification of membrane antigen C33 recognized by monoclonal antibodies inhibitory to human T-cell leukemia virus type 1 (HTLV-1)-induced syncytium formation: altered glycosylation of C33 antigen in HTLV-1-positive T cells. *J Virol.* 1992 Mar;66(3):1394-401.

\*Imai T, Yoshie O: C33 antigen and M38 antigen recognized by monoclonal antibodies inhibitory to syncytium formation by human T cell leukemia virus type 1 are both members of the transmembrane 4 superfamily and associate with each other and with CD4 or CD8 in T cells. *J Immunol.* 1993 Dec 1;151(11):6470-81.

\*Imai T, Kakizaki M, Nishimura M, Yoshie O: Molecular analyses of the association of CD4 with two members of the transmembrane 4 superfamily, CD81 and CD82. *J Immunol.* 1995 Aug 1;155(3):1229-39.

\*Escola JM, Kleijmeer MJ, Stoorvogel W, Griffith JM, Yoshie O, Geuze HJ: Selective enrichment of tetraspan proteins on the internal vesicles of multivesicular endosomes and on exosomes secreted by human B-lymphocytes. *J Biol Chem.* 1998 Aug 7;273(32):20121-7."

\*Stehlíková O, Chovancová J, Tichý B, Krejčí M, Brychtová Y, Panovská A, Francová Skuhrová H, Burčková K, Borský M, Loja T, Mayer J, Pospíšilová S, Doubek M: Detecting minimal residual disease in patients with chronic lymphocytic leukemia using 8-color flow cytometry protocol in routine hematological practice. *Int J Lab Hematol.* 2013 Sep 13. doi: 10.1111/ijlh.12149.

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at [www.exbio.cz](http://www.exbio.cz).

**For laboratory research only, not for drug, diagnostic or other use.**

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic  
Tel: +420 261 090 666 | Fax: +420 261 090 660 | [orders@exbio.cz](mailto:orders@exbio.cz) | [www.exbio.cz](http://www.exbio.cz)