



1F-107-C025

Monoclonal Antibody to Cytokeratin 18 Fluorescein (FITC) conjugated (0.025 mg)

Clone:	DC-10
Isotype:	Mouse IgG1
Specificity:	The antibody DC-10 reacts with Cytokeratin 18 (45 kDa). Cytokeratins are a member of intermediate filaments subfamily represented in epithelial tissues.
Regulatory Status:	RUO
Immunogen:	Human breast carcinoma cell line PMC-42.
Species Reactivity:	Human
Negative Species:	Porcine, Mouse, Rat, Hamster, Bovine, Canine (Dog), Sheep
Preparation:	The purified antibody is conjugated with Fluorescein isothiocyanate (FITC) under optimum conditions. The reagent is free of unconjugated FITC.
Concentration:	0.1 mg/ml
Storage Buffer:	The reagent is provided in stabilizing phosphate buffered saline (PBS) solution containing 15mM sodium azide.
Storage / Stability:	Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light. Do not use after expiration date stamped on vial label.
Usage:	The reagent is designed for Flow Cytometry analysis and Immunofluorescence. Suggested working dilution is 5-10 ug/ml. Indicated dilution is recommended starting point for use of this product. Working concentrations should be determined by the investigator.
Expiration:	See vial label
Lot Number:	See vial label
Background:	Cytokeratins are a subfamily of intermediate filaments and characterized by remarkable biochemical diversity. Cytokeratins are represented in epithelial tissues by at least 20 different polypeptides, molecular weight between 40 kDa and 68 kDa. The individual cytokeratin polypeptides are designated 1 to 20 and divided into the type I (acidic cytokeratins 9-20) and type II (basic to neutral cytokeratins 1-8) families. Cytokeratin 18 belongs to type I family (acidic cytokeratins).
References:	*Lauerova L, Kovarik J, Bartek J, Rejthar A, Vojtesek B: Novel monoclonal antibodies defining epitope of human cytokeratin 18 molecule. <i>Hybridoma</i> . 1988 Oct;7(5):495-504. *Kovarik J, Rejthar A, Lauerova L, Vojtesek B, Bartkova J: Monoclonal antibodies against individual cytokeratins in the detection of metastatic spread. <i>Int J Cancer Suppl</i> . 1988;3:50-5. *Vojtesek B, Staskova Z, Nenutil R, Lauerova L, Kovarik J, Rejthar A, Bartkova J, Bartek J: Monoclonal antibodies recognizing different epitopes of cytokeratin No.18. <i>Folia Biol (Praha)</i> . 1989;35(6):373-82.

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at www.exbio.cz.

For laboratory research only, not for drug, diagnostic or other use.

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic
Tel: +420 261 090 666 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz