

1F-275-T025

Monoclonal Antibody to CD108 Fluorescein (FITC) conjugated (25 tests)

Clone:	MEM-150
Isotype:	Mouse IgM
Specificity:	<p>The antibody MEM-150 reacts with CD108 (JMH blood group antigen), a 80 kDa GPI-anchored glycoprotein expressed on various cell types including erythrocytes, lymphoblasts; at low levels it is present on circulating lymphocytes.</p> <p>HLDA V; WS Code AS S017 HLDA V; WS Code BP BP347 HLDA VI; WS Code BP 401 HLDA VI; WS Code BP 475 HLDA VI; WS Code NL N-L156 HLDA VI; WS Code P PR-65</p>
Regulatory Status:	RUO
Immunogen:	HPB-ALL human T cell line
Species Reactivity:	Human
Preparation:	The purified antibody is conjugated with Fluorescein isothiocyanate (FITC) under optimum conditions. The reagent is free of unconjugated FITC and adjusted for direct use. No reconstitution is necessary.
Storage Buffer:	The reagent is provided in stabilizing Tris buffered saline (TBS) solution containing 15 mM sodium azide.
Storage / Stability:	Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light. Do not use after expiration date stamped on vial label.
Usage:	<p>The reagent is designed for Flow Cytometry analysis of human blood cells using 20 µl reagent / 100 µl of whole blood or 10⁶ cells in a suspension.</p> <p>The content of a vial (0.5 ml) is sufficient for 25 tests.</p>
Expiration:	See vial label
Lot Number:	See vial label
Background:	<p>CD108 (Sema7A) is a GPI-anchored semaphorin family member, which enhances central and peripheral axonal growth and is required for proper axon track formation during embryogenesis. CD108 also regulates osteoclast differentiation and pre-osteoblastic cell migration, and in immune system affects cell proliferation, chemotaxis and cytokine release. On erythrocytes CD108 defines the JMH (John-Milton-Hagen) human blood group. CD108 signalizes through its receptors plexin C1 and beta1 integrins.</p>

For laboratory research only, not for drug, diagnostic or other use.



Antibodies

- References:**
- *Pasterkamp RJ, Peschon JJ, Spriggs MK, Kolodkin AL: Semaphorin 7A promotes axon outgrowth through integrins and MAPKs. *Nature*. 2003 Jul 24;424(6947):398-405.
 - *Delorme G, Saltel F, Bonnelye E, Jurdic P, Machuca-Gayet I: Expression and function of semaphorin 7A in bone cells. *Biol Cell*. 2005 Jul;97(7):589-97.
 - *Pasterkamp RJ, Kolk SM, Hellemons AJ, Kolodkin AL: Expression patterns of semaphorin7A and plexinC1 during rat neural development suggest roles in axon guidance and neuronal migration. *BMC Dev Biol*. 2007 Aug 29;7:98.
 - *Suzuki K, Okuno T, Yamamoto M, Pasterkamp RJ, Takegahara N, Takamatsu H, Kitao T, Takagi J, Rennert PD, Kolodkin AL, Kumanogoh A, Kikutani H: Semaphorin 7A initiates T-cell-mediated inflammatory responses through alpha1beta1 integrin. *Nature*. 2007 Apr 5;446(7136):680-4.
 - *Mudad R, Rao N, Angelisova P, Horejsi V, Telen MJ.: Evidence that CDw108 membrane protein bears the JMH blood group antigen. *Transfusion*. 1995 Jul;35(7):566-70.
 - *Leukocyte Typing V., Schlossman S. et al. (Eds.), Oxford University Press (1995).
 - *Leukocyte Typing VI., Kishimoto T. et al. (Eds.), Garland Publishing Inc. (1997).
 - *Angelisova P, Drbal K, Cerny J, Hilgert I, Horejsi V: Characterization of the human leukocyte GPI-anchored glycoprotein CDw108 and its relation to other similar molecules. *Immunobiology*. 1999 Jun;200(2):234-45.

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at www.exbio.cz.

For laboratory research only, not for drug, diagnostic or other use.

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic
Tel: +420 261 090 666 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz