



1F-297-T100

Monoclonal Antibody to CD34 Fluorescein (FITC) conjugated (100 tests)

Clone:	4H11[APG]
Isotype:	Mouse IgG1
Specificity:	The antibody 4H11[APG] reacts with Class III epitope on CD34 (Mucosialin), a 110-115 kDa monomeric transmembrane phosphoglycoprotein expressed on hematopoietic progenitor cells and on the most pluripotential stem cells; it is gradually lost on progenitor cells. The antibody 4H11[APG] completely blocks binding of Class II antibody QBEnd10 and Class III antibodies BIRMA K3 and 8G12 on KG1a cell line. HLDA VI; WS Code M MA58
Regulatory Status:	RUO
Immunogen:	Permanent human cell line derived from peripheral leucocytes of a patient suffering from chronic myeloid leukaemia.
Species Reactivity:	Human
Preparation:	The purified antibody is conjugated with Fluorescein isothiocyanate (FITC) under optimum conditions. The reagent is free of unconjugated FITC and adjusted for direct use. No reconstitution is necessary.
Storage Buffer:	The reagent is provided in stabilizing phosphate buffered saline (PBS) solution containing 15mM sodium azide.
Storage / Stability:	Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light. Do not use after expiration date stamped on vial label.
Usage:	The reagent is designed for Flow Cytometry analysis of human blood cells using 20 µl reagent / 100 µl of whole blood or 10 ⁶ cells in a suspension. The content of a vial (2 ml) is sufficient for 100 tests.
Expiration:	See vial label
Lot Number:	See vial label
Background:	CD34 is a highly glycosylated monomeric 111-115 kDa surface protein, which is present on many stem cell populations. It is a well established stem cell marker, though its expression on human hematopoietic stem cells is reversible. CD34 probably serves as a surface receptor that undergoes receptor-mediated endocytosis and regulates adhesion, differentiation and proliferation of hematopoietic stem cells and other progenitors. CD34 expression is likely to represent a specific state of hematopoietic development that may have altered adhering properties with expanding and differentiating capabilities in both in vitro and in vivo conditions.

For laboratory research only, not for drug, diagnostic or other use.



Antibodies

- References:**
- *Krauter J, Hartl M, Hambach L, Kohlenberg A, Gunsilius E, Ganser A, Heil G: Receptor-mediated endocytosis of CD34 on hematopoietic cells after stimulation with the monoclonal antibody anti-HPCA-1. *J Hematother Stem Cell Res.* 2001 Dec;10(6):863-71.
 - *Dao MA, Arevalo J, Nolta JA: Reversibility of CD34 expression on human hematopoietic stem cells that retain the capacity for secondary reconstitution. *Blood.* 2003 Jan 1;101(1):112-8.
 - *Gangenahalli GU, Singh VK, Verma YK, Gupta P, Sharma RK, Chandra R, Gulati S, Luthra PM: Three-dimensional structure prediction of the interaction of CD34 with the SH3 domain of Crk-L. *Stem Cells Dev.* 2005 Oct;14(5):470-7.
 - *Gangenahalli GU, Singh VK, Verma YK, Gupta P, Sharma RK, Chandra R, Luthra PM: Hematopoietic stem cell antigen CD34: role in adhesion or homing. *Stem Cells Dev.* 2006 Jun;15(3):305-13.
 - *Leukocyte Typing VI., Kishimoto T. et al. (Eds.), Garland Publishing Inc. (1997).
 - *Elknerová K, Lacinová Z, Soucek J, Marinov I, Stöckbauer P: Growth inhibitory effect of the antibody to hematopoietic stem cell antigen CD34 in leukemic cell lines. *Neoplasma.* 2007;54(4):311-20.

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at www.exbio.cz.

For laboratory research only, not for drug, diagnostic or other use.

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic
Tel: +420 261 090 666 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz