



1F-320-C025

Monoclonal Antibody to IgM Fluorescein (FITC) conjugated (0.025 mg)

Clone:	CH2
Isotype:	Mouse IgG1
Specificity:	The antibody CH2 reacts with Fc fragment of human IgM.
Regulatory Status:	RUO
Immunogen:	Purified human IgM.
Species Reactivity:	Human
Preparation:	The purified antibody is conjugated with Fluorescein isothiocyanate (FITC) under optimum conditions. The reagent is free of unconjugated FITC.
Concentration:	1 mg/ml
Storage Buffer:	Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.4
Storage / Stability:	Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light. Do not use after expiration date stamped on vial label.
Usage:	The reagent is designed for Flow Cytometry analysis. Suggested working dilution is 1:500. Indicated dilution is recommended starting point for use of this product. Working concentrations should be determined by the investigator.
Expiration:	See vial label
Lot Number:	See vial label
Background:	Immunoglobulin M (IgM) is produced as a 900 kDa pentamer, which is an efficient complement binder. This antibody type is produced initially in the immune response and it is the first immunoglobulin class to be synthesized by a fetus or newborn. IgM antibodies do not cross the placenta. IgM concentration in blood is 0.12 g/l and its biological survival (plasma T1/2) is 5 days.
References:	*Franklin EC: Structure and function of immunoglobulins. Acta Endocrinol Suppl (Copenh). 1975;194:77-95. *Fuller JM, Keyser JW: Serum immunoglobulins after surgical operation. Clin Chem. 1975 May;21(6):667-71. *Balogh Z, Merétey K, Falus A, Bozsóky S: Serological abnormalities in juvenile chronic arthritis: a review of 46 cases. Ann Rheum Dis. 1980 Apr;39(2):129-34. *Brinkmann V, Heusser CH: T cell-dependent differentiation of human B cells into IgM, IgG, IgA, or IgE plasma cells: high rate of antibody production by IgE plasma cells, but limited clonal expansion of IgE precursors. Cell Immunol. 1993 Dec;152(2):323-32.

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at www.exbio.cz.

For laboratory research only, not for drug, diagnostic or other use.

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic
Tel: +420 261 090 666 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz