



1F-432-C100

Monoclonal Antibody to HLA-G Fluorescein (FITC) conjugated (0.1 mg)

Clone:	2A12
Isotype:	Mouse IgG1
Specificity:	The antibody 2A12 was generated to a peptide corresponding to C-intron 4-encoded sequence. This antibody does not crossreact with the full-length HLA-G1 isoform and thus allow to distinguish between secreted HLA-G5 and HLA-G6 isoforms from shedded HLA-G1.
Regulatory Status:	RUO
Immunogen:	C-terminal amino acid sequence (22-mer) of soluble HLA-G5 and HLA-G6 proteins coupled to ovalbumin.
Species Reactivity:	Human
Preparation:	The purified antibody is conjugated with Fluorescein isothiocyanate (FITC) under optimum conditions. The reagent is free of unconjugated FITC.
Concentration:	1 mg/ml
Storage Buffer:	Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.4
Storage / Stability:	Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light. Do not use after expiration date stamped on vial label.
Usage:	The reagent is designed for Flow Cytometry analysis. Suggested working dilution is 5 µg/ml. Indicated dilution is recommended starting point for use of this product. Working concentrations should be determined by the investigator.
Expiration:	See vial label
Lot Number:	See vial label
References:	*LeMaoult J, Le Discorde M, Rouas-Freiss N, Moreau P, Menier C, McCluskey J, Carosella ED: Biology and functions of human leukocyte antigen-G in health and sickness. <i>Tissue Antigens</i> . 2003 Oct;62(4):273-84. *White SR, Loisel DA, McConville JF, Stern R, Tu Y, Marroquin BA, Noth I, Ober C: Levels of soluble human leukocyte antigen-G are increased in asthmatic airways. <i>Eur Respir J</i> . 2010 Apr;35(4):925-7. *Larsen MH, Bzorek M, Pass MB, Larsen LG, Nielsen MW, Svendsen SG, Lindhard A, Hviid TV: Human leukocyte antigen-G in the male reproductive system and in seminal plasma. <i>Mol Hum Reprod</i> . 2011 Dec;17(12):727-38. *Poláková K, Železníková T, Russ G: HLA-G5 in the blood of leukemia patients and healthy individuals. <i>Leuk Res</i> . 2013 Feb;37(2):139-45.

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at www.exbio.cz.

For laboratory research only, not for drug, diagnostic or other use.

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic
Tel: +420 261 090 666 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz