

1F-438-C025

Monoclonal Antibody to HLA-G Fluorescein (FITC) conjugated (0.025 mg)

Clone:	01G
Isotype:	Mouse IgG1
Specificity:	The antibody 01G recognizes membrane-bound form of HLA-G (full-length HLA-G1), but not soluble forms. HLA-G belongs to the MHC Class I molecules (MHC Class Ib; nonclassical) and it is expressed on the surface of trophoblast cells.
Regulatory Status:	RUO
Immunogen:	HLA-B27 transgenic mice were immunized with H-2 identical murine cells transfected with and expressing genes encoding HLA-G and human beta2-microglobulin.
Species Reactivity:	Human
Negative Species:	Mouse, Rat
Preparation:	The purified antibody is conjugated with Fluorescein isothiocyanate (FITC) under optimum conditions. The reagent is free of unconjugated FITC.
Concentration:	1 mg/ml
Storage Buffer:	Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.4
Storage / Stability:	Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light. Do not use after expiration date stamped on vial label.
Usage:	The reagent is designed for Flow Cytometry analysis of cells expressing HLA-G molecule on the cell surface. Suggested working dilution is 1:200. Indicated dilution is recommended starting point for use of this product. Working concentrations should be determined by the investigator.
Expiration:	See vial label
Lot Number:	See vial label

For laboratory research only, not for drug, diagnostic or other use.



Antibodies

References:

- *Ishitani A. et al., Protein Expression and Peptide Binding Suggest Unique and Interacting Functional Roles for HLA-E, F and G in Maternal-Placental Immune Recognitions. *The Journal of Immunology* 171, 1376 (2003).
- *Real LM, Cabrera T, Collado A, Jimenez P, Garcia A, Ruiz-Cabello F, Garrido F: Expression of HLA G in human tumors is not a frequent event. *Int J Cancer*. 1999 May 17;81(4):512-8.
- *Real LM, Cabrera T, Canton J, Oliva R, Ruiz-Cabello F, Garrido F: Looking for HLA-G expression in human tumours. *J Reprod Immunol*. 1999 Jul;43(2):263-73.
- *Polakova K, Russ G: Expression of the non-classical HLA-G antigen in tumor cell lines is extremely restricted. *Neoplasma*. 2000;47(6):342-8.
- *Polakova K, Bandzuchova E, Hofmeister V, Weiss EH, Hutter H, Russ G: Binding analysis of HLA-G specific antibodies to hematopoietic cells isolated from leukemia patients. *Neoplasma*. 2003;50(5):331-8.
- *Polakova K, Krcova M, Kuba D, Russ G: Analysis of HLA-G expression in malignant hematopoietic cells from leukemia patients. *Leuk Res*. 2003 Jul;27(7):643-8.
- *Wiendl H, Feger U, Mittelbronn M, Jack C, Schreiner B, Stadelmann C, Antel J, Brueck W, Meyermann R, Bar-Or A, Kieseier BC, Weller M: Expression of the immune-tolerogenic major histocompatibility molecule HLA-G in multiple sclerosis: implications for CNS immunity. *Brain*. 2005 Nov;128(Pt 11):2689-2704.

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at www.exbio.cz.

For laboratory research only, not for drug, diagnostic or other use.

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic
Tel: +420 261 090 666 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz