



1F-482-T025

## Monoclonal Antibody to CD5 Fluorescein (FITC) conjugated (25 tests)

<b>Clone:</b>	CRIS1
<b>Isotype:</b>	Mouse IgG2a
<b>Specificity:</b>	The antibody CRIS1 reacts with the cell surface glycoprotein CD5, a 67kDa single-chain transmembrane glycoprotein expressed on mature T lymphocytes, most of thymocytes and B lymphocytes subset (B-1a lymphocytes). HLDA I; WS Code T 29 HLDA III; WS Code T 530
<b>Regulatory Status:</b>	RUO
<b>Immunogen:</b>	stimulated human leukocytes
<b>Species Reactivity:</b>	Human, Other species Not tested
<b>Preparation:</b>	The purified antibody is conjugated with Fluorescein isothiocyanate (FITC) under optimum conditions. The reagent is free of unconjugated FITC and adjusted for direct use. No reconstitution is necessary.
<b>Storage Buffer:</b>	The reagent is provided in stabilizing phosphate buffered saline (PBS) solution containing 15mM sodium azide.
<b>Storage / Stability:</b>	Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light. Do not use after expiration date stamped on vial label.
<b>Usage:</b>	The reagent is designed for Flow Cytometry analysis of human blood cells using 20 µl reagent / 100 µl of whole blood or 10 <sup>6</sup> cells in a suspension. The content of a vial (0.5 ml) is sufficient for 25 tests.
<b>Expiration:</b>	See vial label
<b>Lot Number:</b>	See vial label
<b>Background:</b>	<p>CD5 antigen (T1; 67 kDa) is a human cell surface T-lymphocyte single-chain transmembrane glycoprotein. CD5 is expressed on all mature T-lymphocytes, most of thymocytes, subset of B-lymphocytes and on many T-cell leukemias and lymphomas. It is a type I membrane glycoprotein whose extracellular region contains three scavenger receptor cysteine-rich (SRCR) domains.</p> <p>The CD5 is a signal transducing molecule whose cytoplasmic tail is devoid of any intrinsic catalytic activity. CD5 modulates signaling through the antigen-specific receptor complex (TCR and BCR). CD5 crosslinking induces extracellular Ca<sup>++</sup> mobilization, tyrosine phosphorylation of intracellular proteins and DAG production. Preliminary evidence shows protein associations with ZAP-70, p56lck, p59fyn, PC-PLC, etc. CD5 may serve as a dual receptor, giving either stimulatory or inhibitory signals depending both on the cell type and development stage. In thymocytes and B1a cells seems to provide inhibitory signals, in peripheral mature T lymphocytes it acts as a costimulatory signal receptor. CD5 is the phenotypic marker of a B cell subpopulation involved in the production of autoreactive antibodies.</p> <p>Disease relevance: CD5 is a phenotypic marker for some B cell lymphoproliferative disorders (B-CLL, Hairy cell leukemia, etc.). The CD5<sup>+</sup> population is expanded in some autoimmune disorders (Rheumatoid Arthritis, etc.). Herpes virus infections induce loss of CD5 expression in the expanded CD8<sup>+</sup> human T cells.</p>

**For laboratory research only, not for drug, diagnostic or other use.**



**Antibodies**

- References:**
- \*Freedman AS, Freeman G, Whitman J, Segil J, Daley J, Levine H, Nadler LM: Expression and regulation of CD5 on in vitro activated human B cells. *Eur J Immunol.* 1989 May;19(5):849-55.
  - Raman C: CD5, an important regulator of lymphocyte selection and immune tolerance. *Immunol Res.* 2002;26(1-3):255-63.
  - \*Leukocyte Typing III., McMichael A. J. et al. (Eds.), Oxford University Press (1987).
  - \*Arrizabalaga P, Mirapeix E, Darnell A, Torras A, Revert L: Cellular immunity analysis using monoclonal antibodies in human glomerulonephritis. *Nephron.* 1989;53(1):41-9.
  - \*Alberola-Ila J, Places L, Cantrell DA, Vives J, Lozano F: Intracellular events involved in CD5-induced human T cell activation and proliferation. *J Immunol.* 1992 Mar 1;148(5):1287-93.
  - \*Guarne A, Bravo J, Calvo J, Lozano F, Vives J, Fita I: Conformation of the hypervariable region L3 without the key proline residue. *Protein Sci.* 1996 Jan;5(1):167-9.

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at [www.exbio.cz](http://www.exbio.cz).

**For laboratory research only, not for drug, diagnostic or other use.**

---

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic  
Tel: +420 261 090 666 | Fax: +420 261 090 660 | [orders@exbio.cz](mailto:orders@exbio.cz) | [www.exbio.cz](http://www.exbio.cz)