

1F-504-T100

Monoclonal Antibody to Kappa light chains Fluorescein (FITC) conjugated (100 tests)

Clone:	A8B5
Isotype:	Mouse IgG1
Specificity:	The antibody A8B5 reacts with kappa light chains (22.5 kDa) of immunoglobulins.
Regulatory Status:	RUO
Species Reactivity:	Human
Negative Species:	Hamster, Sheep, Goat, Rabbit, Guinea pig
Preparation:	The purified antibody is conjugated with Fluorescein isothiocyanate (FITC) under optimum conditions. The reagent is free of unconjugated FITC and adjusted for direct use. No reconstitution is necessary.
Storage Buffer:	The reagent is provided in stabilizing phosphate buffered saline (PBS) solution containing 15mM sodium azide.
Storage / Stability:	Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light. Do not use after expiration date stamped on vial label.
Usage:	The reagent is designed for Flow Cytometry analysis of human blood cells using 20 µl reagent / 100 µl of whole blood or 10 ⁶ cells in a suspension. The content of a vial (2 ml) is sufficient for 100 tests.
Expiration:	See vial label
Lot Number:	See vial label
Background:	Immunoglobulin classes share the same basic four polypeptide chain structure of two heavy chains (five heavy chains types) and two light chains (kappa, lambda; both having a molecular weight of 22.5kDa). Kappa and lambda consist of a variable region and a constant region and can easily be differentiated by the antigenic properties of the constant region. The ratio of kappa to lambda is 70:30.

For laboratory research only, not for drug, diagnostic or other use.

**Antibodies**

- References:**
- *Franklin EC: Structure and function of immunoglobulins. Acta Endocrinol Suppl (Copenh). 1975;194:77-95.
 - *Konig A, Wennemuth G, Soyer HP, Hoffmann R, Happle R, Krause W: Vulvar amyloidosis mimicking giant condylomata acuminata in a patient with multiple myeloma. Eur J Dermatol. 1999 Jan-Feb;9(1):29-31.
 - *Avilés-Salas A, Chanona-Vilchis JG, Candelaria-Hernández MG, Peñalva-Torres Mde L, Cuéllar-Mendoza ME: Presentación inicial de un linfoma plasmablastico como tumor ovárico. Ginecol Obstet Mex. 2012 Jun;80(6):430-4.
 - *Diaz de Domingo NB, Lardo MM, Romero Artaza J, Merelli A, Carbia CD, Sánchez Avalos JC: Aplicacion de tecnicas inmunocitoquimicas para el diagnostico de neoplasias hematologicas. Medicina (B Aires). 1999;59(1):11-6.
 - *Scheeren FA, Diehl SA, Smit LA, Beaumont T, Naspetti M, Bende RJ, Blom B, Karube K, Ohshima K, van Noesel CJ, Spits H: IL-21 is expressed in Hodgkin lymphoma and activates STAT5: evidence that activated STAT5 is required for Hodgkin lymphomagenesis. Blood. 2008 May 1;111(9):4706-15. doi: 10.1182/blood-2007-08-105643
 - *Di Bartolo DL, Hyjek E, Keller S, Guasparri I, Deng H, Sun R, Chadburn A, Knowles DM, Cesarman E: Role of defective Oct-2 and OCA-B expression in immunoglobulin production and Kaposi's sarcoma-associated herpesvirus lytic reactivation in primary effusion lymphoma. J Virol. 2009 May;83(9):4308-15. doi: 10.1128/JVI.02196-08.

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at www.exbio.cz.

For laboratory research only, not for drug, diagnostic or other use.

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic
Tel: +420 261 090 666 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz