



1F-657-C025

Monoclonal Antibody to CD161 (rat) Fluorescein (FITC) conjugated (0.025 mg)

Clone:	10/78
Isotype:	Mouse IgG1
Specificity:	The mouse monoclonal antibody 10/78 recognizes CD161, an approximately 30 kDa type II transmembrane C-type lectin receptor, expressed on the plasma membrane of NK cells, dendritic cells, activated monocytes and a subset of T cells as a disulphide-linked homodimer. A common epitope on rat CD161a and b isoforms is detected.
Regulatory Status:	RUO
Immunogen:	Splenic cells purified from the LEW rat
Species Reactivity:	Rat
Preparation:	The purified antibody is conjugated with Fluorescein isothiocyanate (FITC) under optimum conditions. The reagent is free of unconjugated FITC.
Concentration:	0.5 mg/ml
Storage Buffer:	Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.4
Storage / Stability:	Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light. Do not use after expiration date stamped on vial label.
Usage:	The reagent is designed for Flow Cytometry analysis.
Expiration:	See vial label
Lot Number:	See vial label
Background:	CD161, also known as Nkrp1 (natural killer receptor protein 1) or Klrb1 (killer cell lectin-like receptor subfamily b member 1), is a disulphide-linked homodimeric receptor, which is involved in regulation of NK cell and NKT cell function. It is expressed on rat NK cells, subset of T cells, dendritic cells, and activated monocytes. Although human CD161 is expressed as one isoform, the rat CD161 has three isoforms, referred to as CD161a, b, and c. These proteins contain C-terminal C-type lectin extracellular domain, a transmembrane domain, and N-terminal intracellular domain, which contains ITIM motif, such as CD161b, and displays inhibitory function, or does not contain ITIM motif, thus also not the inhibitory function, such as CD161a.

For laboratory research only, not for drug, diagnostic or other use.



Antibodies

- References:**
- *Kraus E, Lambracht D, Wonigeit K, Hünig T: Negative regulation of rat natural killer cell activity by major histocompatibility complex class I recognition. *Eur J Immunol.* 1996 Nov;26(11):2582-6.
 - *Sedgwick JD, Ford AL, Foulcher E, Airriess R: Central nervous system microglial cell activation and proliferation follows direct interaction with tissue-infiltrating T cell blasts. *J Immunol.* 1998 Jun 1;160(11):5320-30.
 - *Tliba O, Chauvin A, Le Vern Y, Boulard C, Sbille P: Evaluation of the hepatic NK cell response during the early phase of *Fasciola hepatica* infection in rats. *Vet Res.* 2002 May-Jun;33(3):327-32.
 - *May E, Dorris ML, Satumtira N, Iqbal I, Rehman MI, Lightfoot E, Taurog JD: CD8 alpha beta T cells are not essential to the pathogenesis of arthritis or colitis in HLA-B27 transgenic rats. *J Immunol.* 2003 Jan 15;170(2):1099-105.
 - *Stephens LA, Barclay AN, Mason D: Phenotypic characterization of regulatory CD4+CD25+ T cells in rats. *Int Immunol.* 2004 Feb;16(2):365-75.
 - *Chang CJ, Tai KF, Roffler S, Hwang LH: The immunization site of cytokine-secreting tumor cell vaccines influences the trafficking of tumor-specific T lymphocytes and antitumor efficacy against regional tumors. *J Immunol.* 2004 Nov 15;173(10):6025-32.
 - *Teshima R, Nakamura R, Nakamura R, Hachisuka A, Sawada JI, Shibutani M: Effects of exposure to decabromodiphenyl ether on the development of the immune system in rats. *J Health Sci* 2008;54(4):382-389

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at www.exbio.cz.

For laboratory research only, not for drug, diagnostic or other use.

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic
Tel: +420 261 090 666 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz