

1F-665-T025

## Monoclonal Antibody to ZAP-70 Fluorescein (FITC) conjugated (25 tests)

<b>Clone:</b>	1E7.2
<b>Isotype:</b>	Mouse IgG1
<b>Specificity:</b>	<p>The mouse monoclonal antibody 1E7.2 recognizes ZAP-70, a 70 kDa protein tyrosine kinase expressed in T and NK cells.</p> <p>ZAP-70 is a molecule susceptible to degradation. It is recommended to use freshly prepared cell lysates (protease inhibitors are essential) to avoid non-specific staining of degradation products.</p> <p>This product is for research and in vitro experimental use only. It is not to be used for any other commercial purpose. Use of this product to produce products for sale or for therapeutic or drug discovery purposes is prohibited. In order to obtain a license to use this product for commercial purposes, contact The Regents of the University of California.</p>
<b>Regulatory Status:</b>	RUO
<b>Immunogen:</b>	A KLH-coupled peptide corresponding to amino acids 282-307 of human ZAP-70
<b>Species Reactivity:</b>	Human, Mouse
<b>Preparation:</b>	The purified antibody is conjugated with Fluorescein isothiocyanate (FITC) under optimum conditions. The reagent is free of unconjugated FITC and adjusted for direct use. No reconstitution is necessary.
<b>Storage Buffer:</b>	The reagent is provided in stabilizing phosphate buffered saline (PBS) solution containing 15mM sodium azide.
<b>Storage / Stability:</b>	Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light. Do not use after expiration date stamped on vial label.
<b>Usage:</b>	<p>The reagent is designed for Flow Cytometry analysis of human blood cells using 10 µl reagent / 100 µl of Permeabilization Medium (originally 50µl of whole blood; see details on the product page)</p> <p>The content of a vial (0.25 ml) is sufficient for 25 tests.</p>
<b>Expiration:</b>	See vial label
<b>Lot Number:</b>	See vial label

**For laboratory research only, not for drug, diagnostic or other use.**

**Antibodies****Background:**

The ZAP-70 (zeta-associated protein of 70 kDa) tyrosine kinase was identified as a tyrosine phosphoprotein that associates with TCR zeta subunit and undergoes tyrosine phosphorylation following TCR stimulation. ZAP-70 is a Syk family tyrosine kinase primarily expressed in T and NK cells that plays an essential role in signaling through the TCR. TCR-mediated activation of T cells is crucial to the immune response. In humans, ZAP-70 gene mutations resulting in lower ZAP-70 protein expression levels or expression of catalytically inactive ZAP-70 proteins, have been identified. ZAP-70 deficiency results in the absence of mature CD8<sup>+</sup> T cells and the prevention of TCR-mediated activation of CD4<sup>+</sup> T cells, and it can lead to severe combined immunodeficiency.

In patients with chronic lymphocytic leukemia (B-CLL), ZAP-70 expression on B cell was shown to be correlated with disease progression and survival. ZAP-70 contains two N-terminal SH2 domains (Src homology domain 2) and a C-terminal kinase domain. During T cell activation, the binding of ZAP-70 SH2 domains to the phosphorylated zeta subunit on the activated TCR complex causes a colocalization with the Lck tyrosine kinase that phosphorylates ZAP-70 on Tyr493 in the activation loop. ZAP-70 autophosphorylates multiple tyrosines in the region between the SH2 domains and the kinase domain, including the binding sites for additional SH2-containing signaling proteins such as SLP76, LAT, Lck, PLCgamma1, Vav, Shc, Ras-GAP, and Abl. ZAP-70-mediated activation of these downstream effectors leads to the release of intracellular calcium stores, and the transcription of interleukin-2 and other genes important for an immune response.

**References:**

- \*Letestu R, Rawstron A, Ghia P, Villamor N, Boeckx N, Boettcher S, Buhl AM, Duerig J, Ibbotson R, Kroeber A, Langerak A, Le Garff-Tavernier M, Mockridge I, Morilla A, Padmore R, Rassenti L, Ritgen M, Shehata M, Smolewski P, Staib P, Tichioni M, Walker C, Ajchenbaum-Cymbalista F: Evaluation of ZAP-70 expression by flow cytometry in chronic lymphocytic leukemia: A multicentric international harmonization process. *Cytometry B Clin Cytom.* 2006 Jul 15;70(4):309-14.
- \*Del Principe MI, Del Poeta G, Buccisano F, Maurillo L, Venditti A, Zucchetto A, Marini R, Niscola P, Consalvo MA, Mazzone C, Ottaviani L, Panetta P, Bruno A, Bomben R, Suppo G, Degan M, Gattei V, de Fabritius P, Cantonetti M, Lo Coco F, Del Principe D, Amadori S: Clinical significance of ZAP-70 protein expression in B-cell chronic lymphocytic leukemia. *Blood.* 2006 Aug 1;108(3):853-61.
- \*Preobrazhensky SN, Bahler DW: Optimization of flow cytometric measurement of ZAP-70 in chronic lymphocytic leukemia. *Cytometry B Clin Cytom.* 2008 Mar;74(2):118-27.
- \*Gachard N, Salviat A, Boutet C, Arnoulet C, Durrieu F, Lenormand B, Leprêtre S, Olschwang S, Jardin F, Lafage-Pochitaloff M, Penther D, Sainty D, Reminieras L, Feuillard J, Béné MC: Multicenter study of ZAP-70 expression in patients with B-cell chronic lymphocytic leukemia using an optimized flow cytometry method. *Haematologica.* 2008 Feb;93(2):215-23.
- \*Sargent RL, Craig FE, Swerdlow SH: Comparison of Bcl-2, CD38 and ZAP-70 Expression in Chronic Lymphocytic Leukemia. *Int J Clin Exp Pathol.* 2009 Jun 16;2(6):574-82.
- \*Vroblova V, Vrbacky F, Hrudkova M, Jankovicova K, Schmitzova D, Maly J, Krejsek J, Smolej L: Significant change in ZAP-70 expression during the course of chronic lymphocytic leukemia. *Eur J Haematol.* 2010 Jun;84(6):513-7.
- \*Rossi FM, Del Principe MI, Rossi D, Irno Consalvo M, Luciano F, Zucchetto A, Bulian P, Bomben R, Dal Bo M, Fangazio M, Benedetti D, Degan M, Gaidano G, Del Poeta G, Gattei V: Prognostic impact of ZAP-70 expression in chronic lymphocytic leukemia: mean fluorescence intensity T/B ratio versus percentage of positive cells. *J Transl Med.* 2010 Mar 8;8:23.
- \*And many other.

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at [www.exbio.cz](http://www.exbio.cz).

**For laboratory research only, not for drug, diagnostic or other use.**

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic  
Tel: +420 261 090 666 | Fax: +420 261 090 660 | [orders@exbio.cz](mailto:orders@exbio.cz) | [www.exbio.cz](http://www.exbio.cz)