



1F-674-T025

Monoclonal Antibody to Kappa light chains Fluorescein (FITC) conjugated (25 tests)

Clone:	TB28-2
Isotype:	Mouse IgG1
Specificity:	The antibody TB28-2 recognizes human Ig kappa light chains of both secreted and cell surface immunoglobulin. It detects also free kappa light chains.
Regulatory Status:	RUO
Immunogen:	Human IgG-kappa myeloma protein
Species Reactivity:	Human
Preparation:	The purified antibody is conjugated with Fluorescein isothiocyanate (FITC) under optimum conditions. The reagent is free of unconjugated FITC and adjusted for direct use. No reconstitution is necessary.
Storage Buffer:	The reagent is provided in stabilizing phosphate buffered saline (PBS) solution containing 15mM sodium azide.
Storage / Stability:	Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light. Do not use after expiration date stamped on vial label.
Usage:	The reagent is designed for Flow Cytometry analysis of human blood cells using 4 µl reagent / 100 µl of whole blood or 10 ⁶ cells in a suspension. The content of a vial (0.1 ml) is sufficient for 25 tests.
Expiration:	See vial label
Lot Number:	See vial label
Background:	Immunoglobulin classes share the same basic four polypeptide chain structure of two heavy chains (five heavy chains types) and two light chains (kappa, lambda; both having a molecular weight of 22.5kDa). Kappa and lambda consist of a variable region and a constant region and can easily be differentiated by the antigenic properties of the constant region. The ratio of kappa to lambda is 70:30.
References:	*Kiyotaki M, Cooper MD, Bertoli LF, Kearney JF, Kubagawa H: Monoclonal anti-Ig antibodies react with varying proportions of human B lineage cells. <i>J Immunol.</i> 1987 Jun 15;138(12):4150-8. *Nakamura T, Kubagawa H, Cooper MD: Heterogeneity of immunoglobulin-associated molecules on human B cells identified by monoclonal antibodies. <i>Proc Natl Acad Sci U S A.</i> 1992 Sep 15;89(18):8522-6. *Böttcher S, Ritgen M, Buske S, Gesk S, Klapper W, Hoster E, Hiddemann W, Unterhalt M, Dreyling M, Siebert R, Kneba M, Pott C; EU MCL MRD Group: Minimal residual disease detection in mantle cell lymphoma: methods and significance of four-color flow cytometry compared to consensus IGH-polymerase chain reaction at initial staging and for follow-up examinations. <i>Haematologica.</i> 2008 Apr;93(4):551-9. *Karandikar NJ, Aquino DB, McKenna RW, Kroft SH: Transient myeloproliferative disorder and acute myeloid leukemia in Down syndrome. An immunophenotypic analysis. <i>Am J Clin Pathol.</i> 2001 Aug;116(2):204-10. *Jourdan M, Caraux A, De Vos J, Fiol G, Larroque M, Cognot C, Bret C, Duperray C, Hose D, Klein B: An in vitro model of differentiation of memory B cells into plasmablasts and plasma cells including detailed phenotypic and molecular characterization. <i>Blood.</i> 2009 Dec 10;114(25):5173-81.

For laboratory research only, not for drug, diagnostic or other use.



Antibodies

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at www.exbio.cz.

For laboratory research only, not for drug, diagnostic or other use.

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic
Tel: +420 261 090 666 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz