



1F-746-T025

Monoclonal Antibody to CD200 Fluorescein (FITC) conjugated (25 tests)

Clone:	OX-104
Isotype:	Mouse IgG1
Specificity:	The mouse monoclonal antibody OX-104 recognizes CD200, a type-1 glycoprotein of the immunoglobulin superfamily, which is expressed in neurons, B and T cell subsets, keratinocytes, follicular dendritic cells, and ovarian cells. HLDA VII; WS Code 70655
Immunogen:	Human CD200
Species Reactivity:	Human
Preparation:	The purified antibody is conjugated with Fluorescein isothiocyanate (FITC) under optimum conditions. The reagent is free of unconjugated FITC and adjusted for direct use. No reconstitution is necessary.
Storage Buffer:	The reagent is provided in stabilizing phosphate buffered saline (PBS) solution containing 15mM sodium azide.
Storage / Stability:	Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light. Do not use after expiration date stamped on vial label.
Usage:	The reagent is designed for Flow Cytometry analysis of human blood cells using 4 µl reagent / 100 µl of whole blood or 10 ⁶ cells in a suspension. The content of a vial (0.1 ml) is sufficient for 25 tests.
Expiration:	See vial label
Lot Number:	See vial label
Background:	CD200 (also known as OX2 or MRC) is a type-1 membrane glycoprotein, which contains two extracellular immunoglobulin domains, transmembrane domain and cytoplasmic domain. It is expressed by neuronal cells, B and T cell subsets, follicular dendritic cells, keratinocytes, and ovarian cells. The interaction between CD200 and its receptor CD200R results in macrophage activation (IL-6 production), inhibition of mast cell degranulation along with reduced TNF-alpha and IL-13 secretion and overall attenuation of the activation status of lymphocytes. It seems CD200 is also involved in maternal tolerance and its decreased expression in hair follicle correlates with follicular miniaturization.

For laboratory research only, not for drug, diagnostic or other use.

**Antibodies****References:**

- *Sopp P, Werling D, Baldwin C: Sopp P, Werling D, Baldwin C: *Vet Immunol Immunopathol.* 2007 Sep 15;119(1-2):106-14
- *Li Y, Zhao LD, Tong LS, Qian SN, Ren Y, Zhang L, Ding X, Chen Y, Wang YX, Zhang W, Zeng XF, Zhang FC, Tang FL, Zhang X, Ba DN, He W, Cao XT, Lipsky PE: Aberrant CD200/CD200R1 expression and function in systemic lupus erythematosus contributes to abnormal T-cell responsiveness and dendritic cell activity. *Arthritis Res Ther.* 2012 May 23;14(3):R123. doi: 10.1186/ar3853.
- *Shiratori I, Yamaguchi M, Suzukawa M, Yamamoto K, Lanier LL, Saito T, Arase H: Down-regulation of basophil function by human CD200 and human herpesvirus-8 CD200. *J Immunol.* 2005 Oct 1;175(7):4441-9.
- *Colmont CS, Benketah A, Reed SH, Hawk NV, Telford WG, Ohshima M, Udey MC, Yee CL, Vogel JC, Patel GK: CD200-expressing human basal cell carcinoma cells initiate tumor growth. *Proc Natl Acad Sci U S A.* 2013 Jan 22;110(4):1434-9.
- *Foster-Cuevas M, Wright GJ, Puklavec MJ, Brown MH, Barclay AN: Human herpesvirus 8 K14 protein mimics CD200 in down-regulating macrophage activation through CD200 receptor. *J Virol.* 2004 Jul;78(14):7667-76.
- *Meuth SG, Simon OJ, Grimm A, Melzer N, Herrmann AM, Spitzer P, Landgraf P, Wiendl H: CNS inflammation and neuronal degeneration is aggravated by impaired CD200-CD200R-mediated macrophage silencing. *J Neuroimmunol.* 2008 Feb;194(1-2):62-9.
- *Garza LA, Yang CC, Zhao T, Blatt HB, Lee M, He H, Stanton DC, Carrasco L, Spiegel JH, Tobias JW, Cotsarelis G: Bald scalp in men with androgenetic alopecia retains hair follicle stem cells but lacks CD200-rich and CD34-positive hair follicle progenitor cells. *J Clin Invest.* 2011 Feb;121(2):613-22

For laboratory research only, not for drug, diagnostic or other use.