



1P-108-C100

Monoclonal Antibody to Cytokeratin (Pan-reactive) Phycoerythrin (PE) conjugated (0.1 mg)

Clone:	C-11
Isotype:	Mouse IgG1
Specificity:	The antibody C-11 reacts with Cytokeratin peptides 4, 5, 6, 8, 10, 13, 18. Cytokeratins are a member of intermediate filaments subfamily represented in epithelial tissues.
Regulatory Status:	RUO
Immunogen:	Keratin-enriched preparation from human epidermoid carcinoma cell line A431.
Species Reactivity:	Mammalian
Preparation:	The purified antibody is conjugated with R-Phycoerythrin (PE) under optimum conditions. The conjugate is purified by size-exclusion chromatography.
Concentration:	0.1 mg/ml
Storage Buffer:	The reagent is provided in stabilizing phosphate buffered saline (PBS) solution containing 15mM sodium azide.
Storage / Stability:	Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light. Do not use after expiration date stamped on vial label.
Usage:	The reagent is designed for Flow Cytometry analysis. It is recommended to use 10-15 µl of antibody conjugate per 10 ⁶ cells (100 µl cell suspension). Since applications vary, the reagent should be titrated for each particular testing system.
Expiration:	See vial label
Lot Number:	See vial label
Background:	Cytokeratins are a subfamily of intermediate filaments and characterized by remarkable biochemical diversity. Cytokeratins are represented in epithelial tissues by at least 20 different polypeptides, molecular weight between 40 kDa and 68 kDa. The individual cytokeratin polypeptides are designated 1 to 20 and divided into the type I (acidic cytokeratins 9-20) and type II (basic to neutral cytokeratins 1-8) families.
References:	*Kovarik J, Rejthar A, Lauerova L, Vojtesek B, Bartkova J: Monoclonal antibodies against individual cytokeratins in the detection of metastatic spread. <i>Int J Cancer Suppl.</i> 1988;3:50-5. *Vojtesek B, Stasková Z, Nenutil R, Lauerová L, Kovarik J, Rejthar A, Bártková J, Bártek J: Monoclonal antibodies recognizing different epitopes of cytokeratin No.18. <i>Folia Biol (Praha).</i> 1989;35(6):373-82. *Bartek J, Vojtesek B, Staskova Z, Bartkova J, Kerekes Z, Rejthar A, Kovarik J: A series of 14 new monoclonal antibodies to keratins: characterization and value in diagnostic histopathology. <i>J Pathol.</i> 1991 Jul;164(3):215-24. *Hamakawa H, Sumida T, Tanioka H, Sogawa K, Yamada T: Extraction of cytokeratin from the human submandibular gland and its electrophoretic analysis. <i>Res Commun Mol Pathol Pharmacol.</i> 1998 Aug;101(2):115-26. *Broekema M, Harmsen MC, Koerts JA, Petersen AH, van Luyn MJ, Navis G, Popa ER: Determinants of tubular bone marrow-derived cell engraftment after renal ischemia/reperfusion in rats. <i>Kidney Int.</i> 2005 Dec;68(6):2572-81.

For laboratory research only, not for drug, diagnostic or other use.



Antibodies

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at www.exbio.cz.

For laboratory research only, not for drug, diagnostic or other use.

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic
Tel: +420 261 090 666 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz