

1P-161-T025

## Monoclonal Antibody to CD193 Phycoerythrin (PE) conjugated (25 tests)

Clone:	5E8
lsotype:	Mouse IgG2b
Specificity:	The mouse monoclonal antibody 5E8 recognizes CD193 (chemokine receptor 3), an approximately 41 kDa protein expressed above all in eosinophils and basophils.
Regulatory Status:	RUO
Species Reactivity:	Human
Preparation:	The purified antibody is conjugated with R-Phycoerythrin (PE) under optimum conditions. The conjugate is purified by size-exclusion chromatography and adjusted for direct use. No reconstitution is necessary.
Storage Buffer:	The reagent is provided in stabilizing phosphate buffered saline (PBS) solution containing 15mM sodium azide.
Storage / Stability:	Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light. Do not use after expiration date stamped on vial label.
Usage:	The reagent is designed for Flow Cytometry analysis of human blood cells using 10 $\mu$ l reagent / 100 $\mu$ l of whole blood or 10 <sup>6</sup> cells in a suspension. The content of a vial (0.25 ml) is sufficient for 25 tests.
Expiration:	See vial label
Lot Number:	See vial label
Background:	CD193 / CCR3 is a G-protein coupled receptor for several chemokines, namely CCL11 (eotaxin), CCL26 (eotaxin-3), CCL7 (MCP-4), or CCL5 (RANTES). It is highly expressed on eosinophils and basophils, and is also detected in TH1 and TH2 cells, as well as in airway epithelial cells. CD193 is the key eosinophil chemokine receptor responsible for regulation of eosinophil migration and function. This receptor may contribute to the accumulation and activation of eosinophils and other inflammatory cells in the allergic airway. It is also known to be an entry co-receptor for HIV-1.
References:	*Morshed M, Hlushchuk R, Simon D, Walls AF, Obata-Ninomiya K, Karasuyama H, Djonov V, Eggel A, Kaufmann T, Simon HU, Yousefi S: NADPH oxidase-independent formation of extracellular DNA traps by basophils. J Immunol. 2014 Jun 1;192(11):5314-23.

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at www.exbio.cz.

For laboratory research only, not for drug, diagnostic or other use.

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic Tel: +420 261 090 666 | Fax: +420 261 090 660 | orders@exbio.cz | www.exbio.cz