

1P-205-T100

## Monoclonal Antibody to CD6 Phycoerythrin (PE) conjugated (100 tests)

Clone: MEM-98

**Isotype:** Mouse IgG1

Specificity: The MEM-98 antibody reacts with CD6, a 100-130 kDa single chain

transmembrane glycoprotein expressed on T and B lymphocytes subsets,

thymocytes, and acute lymphocytic leukemia cells.

Regulatory Status: RUO

Immunogen: Human CD6 antigen purified by immunoaffinity chromatography from HBP-ALL

cells followed by preparative SDS-PAGE of non-boiled non-reduced sample

(excised piece of gel corresponding to the 100 kDa zone).

Species Reactivity: Human

**Preparation:** The purified antibody is conjugated with R-Phycoerythrin (PE) under optimum

conditions. The conjugate is purified by size-exclusion chromatography and

adjusted for direct use. No reconstitution is necessary.

**Storage Buffer:** The reagent is provided in stabilizing phosphate buffered saline (PBS) solution

containing 15mM sodium azide.

**Storage / Stability:** Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light. Do not

use after expiration date stamped on vial label.

Usage: The reagent is designed for Flow Cytometry analysis of human blood cells using

20 µl reagent / 100 µl of whole blood or 10<sup>6</sup> cells in a suspension.

The content of a vial (2 ml) is sufficient for 100 tests.

**Expiration:** See vial label

**Lot Number:** See vial label

Background: CD6, also known as T12, is a member of the scavenger receptor superfamily found

on T and B cell subsets, thymocytes, and acute lymphocytic leukemia cells (ALL). CD6 interacts with its ligand CD166/ALCAM (activated leukocyte cell adhesion molecule) and serves as a coreceptor for T cell activation and stabilizer of the immunological synapse. CD6-ALCAM mediated cell adhesion is also important for T cell proliferation. CD6 may exert some its functions via association with CD5, probably by fine-tuning CD5 signaling. Ligation of CD6 has antiapoptotic role in

chronic lymphocytic leukemia B cells.



## PRODUCT DATA SHEET

## References:

\*Osorio LM, De Santiago A, Aguilar-Santelises M, Mellstedt H, Jondal M: CD6 ligation modulates the Bcl-2/Bax ratio and protects chronic lymphocytic leukemia B cells from apoptosis induced by anti-IgM. Blood. 1997 Apr 15;89(8):2833-41. \*Zimmerman AW, Joosten B, Torensma R, Parnes JR, van Leeuwen FN, Figdor CG.: Long-term engagement of CD6 and ALCAM is essential for T-cell proliferation induced by dendritic cells. Blood. 2006 Apr 15;107(8):3212-20. Epub 2005 Dec 13. \*Bazil V. et al: Monoclonal antibodies against human leucocyte antigens. III. Antibodies against CD45R, CD6, CD44 and two newly described broadly expressed glycoproteins MEM-53 and MEM-102. Folia Biol. (Praha) 35, 35 (1989). \*Hassan NJ, Simmonds SJ, Clarkson NG, Hanrahan S, Puklavec MJ, Bomb M, Barclay AN, Brown MH: CD6 regulates T-cell responses through activation-dependent recruitment of the positive regulator SLP-76. Mol Cell Biol. 2006 Sep;26(17):6727-38.

\*Castro MA, Oliveira MI, Nunes RJ, Fabre S, Barbosa R, Peixoto A, Brown MH, Parnes JR, Bismuth G, Moreira A, Rocha B, Carmo AM: Extracellular isoforms of CD6 generated by alternative splicing regulate targeting of CD6 to the immunological synapse. J Immunol. 2007 Apr 1;178(7):4351-61.

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at www.exbio.cz.

For laboratory research only, not for drug, diagnostic or other use.