



1P-210-T025

## Monoclonal Antibody to CD11a Phycoerythrin (PE) conjugated (25 tests)

<b>Clone:</b>	MEM-25
<b>Isotype:</b>	Mouse IgG1
<b>Specificity:</b>	The antibody MEM-25 reacts with CD11a (alpha subunit of human LFA-1), a 170-180 kDa type I transmembrane glycoprotein expressed on B and T lymphocytes, monocytes, macrophages, neutrophils, basophils and eosinophils. HLDA IV; WS Code NL 209
<b>Regulatory Status:</b>	RUO
<b>Immunogen:</b>	Leukocytes from a patient suffering from a LGL-type leukaemia.
<b>Species Reactivity:</b>	Human
<b>Preparation:</b>	The purified antibody is conjugated with R-Phycoerythrin (PE) under optimum conditions. The conjugate is purified by size-exclusion chromatography and adjusted for direct use. No reconstitution is necessary.
<b>Storage Buffer:</b>	The reagent is provided in stabilizing phosphate buffered saline (PBS) solution containing 15mM sodium azide.
<b>Storage / Stability:</b>	Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light. Do not use after expiration date stamped on vial label.
<b>Usage:</b>	The reagent is designed for Flow Cytometry analysis of human blood cells using 20 µl reagent / 100 µl of whole blood or 10 <sup>6</sup> cells in a suspension. The content of a vial (0.5 ml) is sufficient for 25 tests.
<b>Expiration:</b>	See vial label
<b>Lot Number:</b>	See vial label
<b>Background:</b>	CD11a (LFA-1 alpha) together with CD18 constitute leukocyte function-associated antigen 1 (LFA-1), the alphaLbeta2 integrin. CD11a is implicated in activation of LFA-1 complex. LFA-1 is expressed on the plasma membrane of leukocytes in a low-affinity conformation. Cell stimulation by chemokines or other signals leads to induction the high-affinity conformation, which supports tight binding of LFA-1 to its ligands, the intercellular adhesion molecules ICAM-1, -2, -3. LFA-1 is thus involved in interaction of various immune cells and in their tissue-specific settlement, but participates also in control of cell differentiation and proliferation and of T-cell effector functions. Blocking of LFA-1 function by specific antibodies or small molecules has become an important therapeutic approach in treatment of multiple inflammatory diseases. For example, humanized anti-LFA-1 antibody Efalizumab (Raptiva) is being used to interfere with T cell migration to sites of inflammation; binding of cholesterol-lowering drug simvastatin to CD11a allosteric site leads to immunomodulation and increase in lymphocytic cholinergic activity.

**For laboratory research only, not for drug, diagnostic or other use.**

**Antibodies****References:**

- \*Sarantos MR, Raychaudhuri S, Lum AF, Staunton DE, Simon SI: Leukocyte function-associated antigen 1-mediated adhesion stability is dynamically regulated through affinity and valency during bond formation with intercellular adhesion molecule-1. *J Biol Chem.* 2005 Aug 5;280(31):28290-8.
- \*Giblin PA, Lemieux RM: LFA-1 as a key regulator of immune function: approaches toward the development of LFA-1-based therapeutics. *Curr Pharm Des.* 2006;12(22):2771-95.
- \*Kellersch B, Kolanus W: Membrane-proximal signaling events in beta-2 integrin activation. *Results Probl Cell Differ.* 2006;43:245-57.
- \*Fujii T, Takada-Takatori Y, Kawashima K: Roles played by lymphocyte function-associated antigen-1 in the regulation of lymphocytic cholinergic activity. *Life Sci.* 2007 May 30;80(24-25):2320-4.
- \*Leukocyte Typing IV., Knapp W. et al. (Eds.), Oxford University Press (1989).
- \*Bazil V, Stefanova I, Hilgert I, Kristofova H, Vanek S, Horejsi V.: Monoclonal antibodies against human leucocyte antigens. IV. Antibodies against subunits of the LFA-1 (CD11a/CD18) leucocyte-adhesion glycoprotein. *Folia Biol (Praha).* 1990;36(1):41-50.
- \*Otonello L, Epstein AL, Mancini M, Dapino P, Dallegri F: Monoclonal LYM-1 antibody-dependent cytotoxicity by human neutrophils exposed to GM-CSF: auto-regulation of target cell attack by cathepsin G. *J Leukoc Biol.* 2004 Jan;75(1):99-105.
- \*Simon SI, Cherapanov V, Nadra I, Waddell TK, Seo SM, Wang Q, Doerschuk CM, Downey GP: Signaling functions of L-selectin in neutrophils: alterations in the cytoskeleton and colocalization with CD18. *J Immunol.* 1999 Sep 1;163(5):2891-901.
- \*Hajshengallis G, Martin M, Sojar HT, Sharma A, Schifferle RE, DeNardin E, Russell MW, Genco RJ: Dependence of bacterial protein adhesins on toll-like receptors for proinflammatory cytokine induction. *Clin Diagn Lab Immunol.* 2002 Mar;9(2):403-11.
- \*Wang JH, Kwas C, Wu L: Intercellular adhesion molecule 1 (ICAM-1), but not ICAM-2 and -3, is important for dendritic cell-mediated human immunodeficiency virus type 1 transmission. *J Virol.* 2009 May;83(9):4195-204.
- \*Garnotel R, Rittié L, Poitevin S, Monboisse JC, Nguyen P, Potron G, Maquart FX, Randoux A, Gillery P: Human blood monocytes interact with type I collagen through alpha x beta 2 integrin (CD11c-CD18, gp150-95). *J Immunol.* 2000 Jun 1;164(11):5928-34.
- \*Mathison RD, Befus AD, Davison JS, Woodman RC: Modulation of neutrophil function by the tripeptide feG. *BMC Immunol.* 2003 Mar 4;4:3. Epub 2003 Mar 4.
- \*Otonello L, Epstein AL, Dapino P, Barbera P, Morone P, Dallegri F: Monoclonal Lym-1 antibody-dependent cytotoxicity by neutrophils exposed to granulocyte-macrophage colony-stimulating factor: intervention of Fc gamma RII (CD32), CD11b-CD18 integrins, and CD66b glycoproteins. *Blood.* 1999 May 15;93(10):3505-11.
- \*Aubert M, Yoon M, Sloan DD, Spear PG, Jerome KR: The virological synapse facilitates herpes simplex virus entry into T cells. *J Virol.* 2009 Jun;83(12):6171-83.
- \*Osicka R, Osickova A, Hasan S, Bumba L, Cerny J, Sebo P: Bordetella adenylate cyclase toxin is a unique ligand of the integrin complement receptor 3. *Elife.* 2015 Dec 9;4:e10766.
- \*And many other.

Unless indicated otherwise, all products are For Research Use Only and not for diagnostic or therapeutic use. Not for resale or transfer either as a stand-alone product or as a component of another product without written consent of EXBIO. EXBIO will not be held responsible for patent infringement or other violations that may occur with the use of our products. All orders are accepted subject to EXBIO's term and conditions which are available at [www.exbio.cz](http://www.exbio.cz).

**For laboratory research only, not for drug, diagnostic or other use.**

EXBIO Praha | Nad Safinou II 341 | 252 50 Vestec u Prahy | Czech Republic  
Tel: +420 261 090 666 | Fax: +420 261 090 660 | [orders@exbio.cz](mailto:orders@exbio.cz) | [www.exbio.cz](http://www.exbio.cz)